

RECLAMATION

Managing Water in the West

2016 Monitoring Report for the Los Lunas Habitat Restoration Project

Middle Rio Grande Project, New Mexico



U.S. Department of the Interior
Bureau of Reclamation
Technical Service Center
Water, Environmental and Ecosystems Division
Fisheries and Wildlife Resources Group
Denver, Colorado

March 2017

Mission Statements

The U.S. Department of the Interior protects America's natural resources and heritage, honors our cultures and tribal communities, and supplies the energy to power our future.

The mission of the Bureau of Reclamation is to manage, develop, and protect water and related resources in an environmentally and economically sound manner in the interest of the American public.

2016 Monitoring Report for the Los Lunas Habitat Restoration Project

Middle Rio Grande Project, New Mexico

Prepared by:

Rebecca Siegle
Natural Resources Specialist

Darrell Ahlers
Wildlife Biologist

Prepared for:

U.S. Army Corps of Engineers
Albuquerque District

U.S. Department of the Interior
Bureau of Reclamation
Albuquerque Area Office

Cover Photo: Looking south along the “root wad berm”.



U.S. Department of the Interior
Bureau of Reclamation
Technical Service Center
Water, Environmental, and Ecosystems Division
Fisheries and Wildlife Resources Group
Denver, Colorado

March 2017

Contents

	<i>Page</i>
Introduction.....	1
Project Background.....	2
Methods.....	5
Avian Monitoring.....	5
Point Counts.....	5
Cleared/Overbank Area	5
Burned Area	5
Willow Flycatcher and Yellow-billed Cuckoo Surveys	7
Vegetation Monitoring.....	7
Vegetation Transects.....	7
Vegetation Quantification Plots.....	9
Groundwater Monitoring	13
Photo Stations	13
Results	14
Avian Monitoring.....	14
Point Counts.....	14
Cleared/Overbank Area	14
Burned Area	19
Comparisons between Monitoring Areas.....	22
Willow Flycatcher and Yellow-billed Cuckoo Surveys	25
Vegetation Monitoring.....	28
Vegetation Transects.....	28
Vegetation Quantification Plots.....	33
Groundwater Monitoring	36
Monthly Well Monitoring.....	36
Data Logger Well Monitoring	37
Photo Stations	38
Discussion.....	39
Avian Monitoring.....	39
Point Counts.....	39
Cleared/Overbank Area	39
Burned Area	41
Willow Flycatcher and Yellow-billed Cuckoo Surveys	41
Vegetation Monitoring.....	42
Vegetation Transects.....	42
Vegetation Quantification Plots.....	44
Groundwater Monitoring	46
Photo Stations	47
Conclusion and Recommendations	48
Avian Monitoring.....	48
Conclusions.....	48
Recommendations.....	49
Vegetation Monitoring.....	50
Conclusions.....	50

Recommendations	51
Groundwater Monitoring	52
Conclusions	52
Recommendations	52
Photo Stations	52
Conclusions	52
Recommendations	52
Literature Cited	1

Appendices

- Appendix A** - Waypoint Locations for Avian Point Counts, Vegetation Transects, Groundwater Monitoring Wells, and Photo Stations
- Appendix B** - Bird Species Detected During Point Counts and Associated Habitat Guilds
- Appendix C** - Relative Abundance of Individual Bird Species by Area
- Appendix D** - Avian Abundance by Species Guilds
- Appendix E** - Southwestern Willow Flycatcher Survey Forms and Maps 2016
- Appendix F** - Plant List and Total Percent Cover of Plants Detected in the Understory Layer by Individual Species, Life-form, and Cover Type 2003 to 2016
- Appendix G** - Groundwater Monitoring Wells; Monthly Data June 2003 – October 2010
- Appendix H** - Groundwater Monitoring Wells; HOBO Water Level Logger Data June 2011– September 2016
- Appendix I** - Photo Stations 2003 - 2016

Tables

	<i>Page</i>
Table 1. P and R values for simple linear regression analysis between year and relative abundance by guild in the Cleared/Overbank Area.	18
Table 2. P and R values for simple linear regression analysis between year and relative abundance by guild in the Burned Area.....	22
Table 3. Statistical comparisons of relative abundance between areas by year and guild.	24
Table 4. Total percent cover and average height of woody overstory species (>1 m) from 2007 to 2016.	31
Table 5. Proportion of native and introduced species in the understory and overstory layers by year.	32
Table 6. Summary of center plot shrub and tree stem count data gathered at SWFL nest sites (2004 to 2006) and LLRS sites (2015 and 2016)..	34
Table 7. Summary of point-centered quarter and canopy cover data from SWFL nest sites (2004 to 2006) and LLRS sites (2015 and 2016)..	35

Figures	<i>Page</i>
Figure 1. Location of the Los Lunas Restoration Site (LLRS) project area.....	3
Figure 2. Cleared/Overbank and Burned Area point count locations at LLRS.....	6
Figure 3. Vegetation transect, well, and photo station locations at LLRS	8
Figure 4. Locations of vegetation quantification plots in the Burned Area (2016) and in the Cleared/Overbank Area (2015 and 2016) within the LLRS. ...	10
Figure 5. Vegetation quantification plot layout.....	12
Figure 6. Typical SWFL habitat showing two layers of vegetation: shrub and canopy.....	12
Figure 7. Shade plot of the most abundant species detected in the Cleared/ Overbank Area by sample and year	15
Figure 8. MDS ordination of 14 years of species abundance data based on Bray- Curtis similarities within the Cleared/Overbank Area.	16
Figure 9. Relative abundance by species guilds in the Cleared/Overbank Area over time.	17
Figure 10. Species richness by species guilds in the Cleared/Overbank Area over time.	17
Figure 11. Linear trend in average number of mid-story birds per point in relation to year (2003 to 2016) in the Cleared/Overbank Area.....	18
Figure 12. Shade plot of the most abundant species detected in the Burned Area by sample and year.....	20
Figure 13. MDS ordination of 11 years of species abundance data based on Bray- Curtis similarities within the Burned Area	21
Figure 14. Relative abundance by species guilds in the Burned Area over time. .	21
Figure 15. Species richness by species guilds in the Burned Area over time.	22
Figure 16. MDS ordination based on 14 years of square root transformed species abundance data and Bray-Curtis similarities for both the Cleared/Overbank and Burned Areas.	23
Figure 17. Trendlines and R^2 values for relative abundance over time in the Cleared/Overbank Area (2003-2015) and Burned Area (2003, 2004, 2007- 2016).	26
Figure 18. Trendlines and R^2 values for species richness over time in the Cleared/Overbank Area (2003-2015) and Burned Area (2003, 2004, 2007- 2016).	26
Figure 19. WIFL detections and habitat suitability in the vicinity of LLRS within the Belen survey site.	27
Figure 20. Plant species richness from 2003 to 2016.....	28
Figure 21. Statistical results analyzing total vegetation cover over time for various parameters.	29
Figure 22. Relative percent cover of life-forms in the understory layer from 2003 to 2016.	30
Figure 23. MDS ordination of 13 years of plant species cover data based on Bray- Curtis similarities	33

Figures (cont'd)

Page

Figure 24. Hydrologic year (October – September) average discharge (cfs) in the Rio Grande at San Acacia (2002-2007) and at Bosque Farms (2008-2015), and the average total percent plant cover in transects at the LLRS, New Mexico.	38
Figure 25. Diurnal fluctuation (ft) within Well M2 and average discharge (cfs) in the Rio Grande at Bosque Farms, New Mexico from September 2013 through August 2014.....	38
Figure 26. Development of vegetation at the LLRS as seen in 2002 immediately after the site was cleared (left) and in 2014 (right).	40
Figure 27. Evidence of <i>Diorhabda</i> , a biological control beetle released in the Southwest to manage saltcedar, was first observed in the LLRS in 2014.	43
Figure 28. Example of sparse leaf cover on a cottonwood, September 2015, LLRS.	48

Introduction

Riparian cottonwood (*Populus* spp.) and willow (*Salix* spp.) forests are an important ecosystem in the Southwestern United States, providing fish and wildlife habitat, biodiversity, and watershed protection (Hultine et al. 2010). Native riparian habitat is used by a wide range of species and in the southwest about 60 percent of all vertebrates species and 70 percent of all threatened and endangered species are riparian obligates (Poff et al. 2012). Along the Middle Rio Grande in central New Mexico, the endangered southwestern willow flycatcher (*Empidonax traillii extimus*; SWFL) and the threatened western yellow-billed cuckoo (*Coccyzus americanus*; YBCU) are species of particular concern that are dependent on riparian habitat. The destruction of riparian habitats has caused severe declines in these populations, which exist only in fragmented and scattered locations throughout their historic range (USFWS 1997, USFWS 2013).

Native riparian communities, although once abundant, are declining and now comprise <2 percent of the land area in the west (Sprenger 1999, Poff et al. 2012). Declines and degradation of native riparian habitat have been associated with a number of activities that have resulted in hydrologic changes. In the Middle Rio Grande, reservoir construction, regulation of surface flow, groundwater pumping, and water diversions have interfered with hydrological processes such as overbank flooding, floodplain scouring, and sediment deposition within floodplains (Sprenger 1999). These flood control structures and flow management regimes have prevented natural flooding necessary for cottonwood and willow regeneration (Dreesen et al. 2002) and have also led to sections of the riparian forest (“bosque”) being less hydrologically connected to the river than they were in the past, lowering the water table (Cartron et al. 2008). The reproductive biology of cottonwood and willow is strongly tied to fluvial processes (Stromberg 1993). In desert riparian areas, seedling establishment is dependent on late winter and early spring flood flows to deposit moist alluvium on sediment bars during the short period in early spring when native seeds are dispersed (Sprenger 1999, Muldavin et al. 2015). Seeds, which are only viable for several weeks, are reliant upon slowly receding flood flows and water tables so seedling roots can stay in contact with adequate soil moisture. Mature plants often become isolated on high floodplains some distance from the active channel, but continue to remain hydrologically dependent on a shallow riparian water table (Stromberg 1993). Mature tree growth and maintenance depends on groundwater remaining above a depth of about 10 feet (ft) in the bosque (Cartron et al. 2008). For the establishment and development of younger age classes (those typically occupied by SWFLs) the groundwater levels must remain much higher – perhaps less than 5 ft based on data collected in association with the Bosque del Apache and Elephant Butte Sediment Plug Studies conducted on the Middle Rio Grande (Siegle et al. 2015a, Siegle et al. 2015b).

In addition, large areas of the Middle Rio Grande that were historically cottonwood forests have been invaded by exotic woody species, primarily saltcedar (*Tamarix* spp.). Saltcedar, like cottonwood and willow, is dependent upon moist, bare substrates created by receding flood flows for initial germination and survival (Sprenger 1999). Unlike

native species, however, saltcedar disperses seed throughout the growing season allowing greater opportunity to establish than native species. The establishment of exotics, along with a predominately dry floodplain that lacks scouring floods and slows decomposition, have magnified the potential of severe wildland fires because of the massive fuel loads produced (Dreesen et al. 2002, Cartron et al. 2008).

In April of 2000, an area of the bosque near Los Lunas, New Mexico suffered a severe fire that destroyed virtually all of the aboveground vegetation. This area thus presented a unique opportunity for native riparian forest restoration and was designated as the Los Lunas Habitat Restoration Project.

Project Background

Historically, the Los Lunas Habitat Restoration Project fulfilled requirements in one of eight reaches in which habitat restoration was to be conducted in accordance with Element J of the Reasonable and Prudent Alternative (RPA) within the June 2001 Biological Opinion (BO) issued by the U.S. Fish and Wildlife Service (USFWS 2001). Following the fire, the Los Lunas Restoration Site (LLRS) was selected as the first BO restoration area (Figure 1). The U.S. Bureau of Reclamation (Reclamation) Albuquerque Area Office and the U.S. Army Corps of Engineers Albuquerque District have acted as joint lead federal agencies on this project, and the Middle Rio Grande Conservancy District is the primary non-federal cooperator.

The primary objectives of the restoration project were to improve habitat conditions for the Rio Grande silvery minnow (*Hybognathus amarus*; minnow) and SWFL such that, in combination with other elements of the RPA, continued jeopardy to the two species could be avoided.

The design goals were to generate inundation of the project area at flows of greater than or equal to 2,500 cubic ft/second (cfs). For flows below 2,500 cfs, a variety of substrate elevations was integrated into the project design to allow for the inundation of certain regions at lower river stages. This included features such as a network of variable depth side and transverse channels designed to aid in minnow egg retention and provide shallow water/low velocity rearing habitat. In addition, the increased inundation frequency would begin the process of post-fire regeneration of high-value terrestrial habitats in portions within and adjacent to the restoration area to support the recovery of the SWFL.

In April 2002, the initial phase of work began by removing approximately 1,400 jetty jacks and establishing access routes and a staging area. When construction was initiated, the site was largely dominated by thick stands of herbaceous and exotic regrowth. Vegetation was cleared and mulched within the overbank area, access roads, staging area, and disturbance areas next to the levee and root-wad berm. With the removal of jetty jacks completed, crews from Reclamation's Socorro Field Office began clearing, surveying, and excavating the flood plain. Specific areas within the site were revegetated using seed, potted shrubs, or cottonwood and willow poles.




Figure 1. Location of the Los Lunas Restoration Site (LLRS) project area.

Properly functioning riparian areas serve key roles in providing fish and wildlife habitat and preserving water quality and supply. Factors such as water table depth and fluctuation, soil texture, soil salinity, and browsing pressure from livestock and wildlife determine the success of restoration in creating a functioning riparian area (Dreesen et al. 2002). Reclamation's Technical Service Center (TSC) in Denver, Colorado has conducted avian, vegetation, and groundwater monitoring at LLRS since 2003. Although requirements of the BO have been met, this study is being continued to provide information for an adaptive management approach to creating and monitoring potential SWFL habitat. The YBCU was added to the federal Threatened and Endangered Species list after initiation of the LLRS project and therefore was not a primary focus for habitat restoration or this study. The species was addressed to a limited extent in this report, however. Objectives of annual monitoring efforts are to:

- determine the success of restoration at the LLRS in establishing a productive cottonwood/willow riparian community, as well as characterizing factors that may have influenced the outcome;
- assess SWFL habitat suitability/sustainability and identify those variables which contribute to the development of SWFL habitat;
- establish a potential timeframe in which a restored site develops into suitable SWFL habitat under local environmental conditions; and
- provide data for the adaptive management of future restoration efforts in the Middle Rio Grande

Methods

This comprehensive study is comprised of various types of monitoring which include avian point counts and SWFL and YBCU surveys, vegetation transects and quantification plots, groundwater wells, and photo stations. Methods used for each type of monitoring are described below.

Avian Monitoring

Point Counts

Avian monitoring included 5-minute, 50-meter (m) fixed-radius point counts that were conducted 3 times/year during the peak breeding season (late-May to early-July). Point counts took place within two areas that were monitored over a 14-year study period from 2003 to 2016 (waypoint locations are listed in Appendix A). These areas – the Cleared/Overbank and Burned Areas – were located within the LLRS and are separated by a root-wad berm constructed during restoration activities. Only the Cleared/ Overbank Area was monitored for the duration of the study. Point counts were conducted in the Burned Area in 2003, 2004, and 2007 to 2016. The Cleared/Overbank and Burned Areas are described below:

Cleared/Overbank Area

This restoration area, adjacent to the active river channel, was cleared and excavated to allow overbank flooding with regrowth comprised of primarily native and mixed vegetation. Monitoring was conducted at eight points from 2003 to 2006; points at this site were relocated and increased to 12 in 2007 so that: a) the points were more evenly distributed over the area; and b) all areas had the same sample size (Figure 2).

Burned Area

This cottonwood gallery, burned in 2000 and adjacent to the Cleared/Overbank Area, experienced regrowth of mixed vegetation. Point counts were conducted in 2003 and 2004, and after a two year hiatus, monitoring was resumed in 2007 to provide a comparison site. Counts were conducted at seventeen points within this site in 2003 and 2004; points were relocated and decreased to 12 in 2007 so that: a) the points were more evenly distributed and were all within the restoration area; and b) all areas had the same sample size (Figure 2).

Data from the 14 years of monitoring were analyzed to evaluate trends in relative abundance of pooled species guilds over time and statistical comparisons were made between areas. Pooled species guilds were categorized based predominately on nesting habitat and included canopy, cavity, dense shrub, edge, ground shrub, mid-story, open, and water birds. Migrants were also documented but were not included in statistical




Figure 2. Cleared/Overbank and Burned Area point count locations at LLRS (NAIP 2014 natural color photography). A root wad berm separates the 2 areas.

analysis. The table in Appendix B shows the groupings of individual bird species into guilds for analysis purposes as well as scientific names and codes of the bird species. This table serves as a reference for scientific names throughout the report.

Statgraphics statistical software was used to conduct simple linear regressions to test for significant relationships between the abundance of birds and year (*i.e.*, time; Nur et al. 1999). To compare bird abundance between areas by year, the Student's t-test was used for normally distributed data and the Mann-Whitney nonparametric test of medians was used for data that were not normally distributed. Primer-e statistical software was used to generate Multi-dimensional Scaling (MDS) configurations which were used to examine species composition over time and between plots. MDS ordination ranks species similarities and the associated configuration can be interpreted in terms of relative similarity of samples to each other (Clarke et al. 2014).

Willow Flycatcher and Yellow-billed Cuckoo Surveys

Three presence/absence surveys were conducted per year for the endangered SWFL within the LLRS from 2004 through 2016 in accordance with Sogge et al. (2010). Additional surveys were conducted within the same period on both sides of the river in adjacent sections. The project site falls within the BL-25 survey site, which is within the Belen reach between the Los Lunas and Belen bridges. These surveys were part of Reclamation's annual SWFL monitoring program conducted at selected sites along the Rio Grande from Bandelier National Monument to Elephant Butte Reservoir (Moore and Ahlers 2016). Surveys included all willow flycatchers (WIFLs; *Empidonax traillii* spp.) but the subspecies of interest is the southwestern willow flycatcher (SWFL; *Empidonax traillii extimus*). All migrants were considered WIFLs while all resident territories were considered SWFLs.

Four presence/absence surveys were conducted annually for the threatened YBCU in accordance with Halterman et al. (2015). Formal surveys along the Middle Rio Grande began in 2009 but were not initiated within the Belen reach until 2014.

Vegetation Monitoring

Vegetation Transects

Twelve 50-m permanent transects were established at the LLRS between the root-wad berm and the river (the site referred to as the Cleared/Overbank Area in avian monitoring) to document the natural establishment of vegetation in this area (waypoint locations are listed in Appendix A). The area where transects were placed was not revegetated using seed or potted shrubs as were some areas within the restoration site. All transects were evenly distributed in the disturbed area and were oriented perpendicular to the river (Figure 3).

Cover and species composition were measured every 0.5 m along the 50-m transect. For understory measurements, the point-intercept method was used, which entailed recording




Figure 3. Vegetation transect, well, and photo station locations at LLRS (NAIP 2014 natural photography).

the first “hit” for herbaceous plant species and for woody species under 1 m tall. If a plant was not intercepted, then bare soil or litter was recorded. As of 2007, the line-intercept method was used for measuring overstory cover. Canopy cover was measured along each transect by noting the point along the tape where the canopy began and the point at which it ended for each woody species over a meter tall. Because species overlapped in some cases, the sum of the cover for all species did not necessarily reflect the actual percentage of overstory cover along the tape. The percentage of the tape covered by overstory was also calculated. The height of the tallest vegetation within each continuous stretch of the same species was measured.

The methodology used for cover measurements was revised in 2007 to include a separate overstory measure (woody species > 1 m in height). Prior to 2007, the method used to collect understory cover was applied to all vegetation cover measurements, so that if a woody species was intercepted first, then this species was recorded as understory. As vegetation grew in height, the original methodology did not account for overstory as a separate layer, and understory vegetation cover was not fully captured. This phenomenon was first noticed in 2006; therefore understory shrub data from that year is probably more comparable to overstory data from 2007 to 2016. Data were collected between mid-August and mid-September from 2003 through 2016.

Data from the 14 years of monitoring were compared to evaluate any statistically significant changes within vegetation types over time. General linear model (GLM) analysis was applied to test for relationships between total cover and year, while Tukey’s honest significant difference (HSD) procedure was used as a multiple comparison test to evaluate statistically significant differences between years ($\alpha=0.05$) utilizing StatGraphics statistical software. The Tukey’s HSD analysis is a post-test to the GLM and provides a more focused analysis of individual years. Primer-e statistical software was used to generate MDS configurations to examine changes in plant species composition over time.

Total percent cover (i.e., actual cover estimate) was statistically analyzed for understory and overstory vegetation. Relative percent cover was determined for understory life-forms (i.e., native or introduced shrubs <1 m in height, grasses and grass-like species, and forbs). Relative cover is cover of a species or life-form expressed as a percentage of total vegetation.

Vegetation Quantification Plots

Between 2004 and 2006, Reclamation gathered and analyzed vegetation data from 112 SWFL nest sites within the Middle Rio Grande. Results of this study are presented in *Vegetation Quantification of Southwestern Willow Flycatcher Nest Sites* (Moore 2007). In an effort to assess the suitability of developing habitat for breeding SWFLs within LLRS, Reclamation gathered similar vegetation data in 2015 and 2016 at sites that appeared suitable for breeding SWFLs but were currently unoccupied (Figure 4). Three plots were measured within the Cleared/Overbank Area in 2015 and again in 2016. Three plots were added to the study and measured within the Burned Area in 2016. LLRS vegetation quantification data was compared to nest site data presented in Moore (2007).




Figure 4. Locations of vegetation quantification plots in the Burned Area (2016) and in the Cleared/Overbank Area (2015 and 2016) within the LLRS (NAIP 2014 natural color photography).

Most of the data collected in association with the 112 nests represents habitat of exceptional quality for SWFL breeding that was located in the delta of Elephant Butte Reservoir. These habitat conditions may not be achievable in the Los Lunas area, which is approximately 100 miles upstream of the delta and experiences entirely different hydrological conditions and is populated by different plant species. To provide a representative comparison for the LLRS, 22 nests from the Sevilleta/La Joya, Bosque del Apache, and Tiffany Reaches – which have similar conditions - were analyzed separately and also used as comparison data.

Methods were adapted from BBIRD protocol (Martin et al. 1997), similar studies conducted by the New Mexico Natural Heritage Program along the Rio Grande (DeRagon et al. 1995, Ahlers and White 1997, Stoleson and Finch 1999), and University of New Mexico (Peter Stacey, pers. comm.).

Vegetation and habitat data were collected within an 11.35-m radius plot (0.04 hectare (ha) BBIRD-type plot) centered below the selected suitable nest substrate (Figure 5). All trees within the center plot were tallied by species. Stems were considered trees when diameter at breast height (DBH) was greater than 5 centimeters (cm). Average stem density, species and size class composition, and percentage of dead trees were computed for these plots. Trees were divided into three DBH classes: Class I consisted of trees 5 cm to 10 cm DBH, Class II consisted of trees 10 cm to 20 cm, and Class III consisted of trees greater than 20 cm.

Shrubs were measured in four 1 x 4 m shrub plots located at random distances less than 7.35 m from the plot center along each of four radii in cardinal directions. Shrub stems were defined as having a DBH between 0.5 cm and 5 cm. All shrub stems within each shrub plot were counted by species. Stem densities, species composition, and percentage of dead were computed. It should be noted that all stems encountered at breast height within the 1 x 4 shrub plots were counted, not necessarily just those that were rooted. Therefore, measurements do not reflect actual stem densities but provide relative comparisons over time.

Three additional subplots, each with a 5 m radius, were established adjacent to each center plot (Figure 5). Measurements within each quarter of the center plot and of the three smaller subplots were taken for plants in 2 layers: shrub and canopy (Figure 6). Point-centered data included DBH, crown width, and height for each of the 2 layers. Canopy cover visual estimates were also made within each of three canopy layers (0 to 3 m, 3 to 6 m, and >6 m). Estimates were made using a Daubenmire ranking of 0 to 6 where 0 = 0 percent cover, 1 = 1 to 10 percent, 2 = 11 to 25 percent, 3 = 26 to 50 percent, 4 = 51 to 75 percent, 5 = 76 to 90 percent, and 6 = greater than 90 percent cover (the midpoint for each of these ranges was used for analysis purposes. If a subplot fell partially or entirely within an area designated as non-habitat for SWFLs (in this case the river channel), it was excluded from measurements. For center plots, the quarter of the plot (as measured from each cardinal direction) that fell in non-habitat, such as open water, was excluded from data collection.




Figure 5. Vegetation quantification plot layout.




Figure 6. Typical SWFL habitat showing two layers of vegetation: shrub and canopy.

In order to compare the LLRS assessment sites to SWFL nest sites, each dataset was pooled separately and mean values were compared. If LLRS mean values were within 0.5 standard deviations of means calculated in the original study, these parameters were considered suitable for nesting SWFLs.

Groundwater Monitoring

Eleven groundwater monitoring wells were installed along 3 transects running perpendicular to the river: 4 wells on the northern end of the site, 4 in the center, and 3 on the southern end (Figure 3; waypoint locations are listed in Appendix A). All wells were installed using the Army Corps of Engineers (2000) methodology. Wells averaged 5.0 ft in depth, with the groundwater depth at a range of 2.0 to 4.0 ft below the surface at the time of installation. Eight wells were installed in June 2003 and the remaining westernmost three were installed July 2004. The eleven wells were manually monitored every month from date of installment to 2010.

In June 2011, HOBO Water Level Loggers were installed in 9 of the wells and hydrologic measurements were discontinued in 2 of the westernmost wells. Loggers were attached to the well cap via a braided stainless steel wire and programmed to collect readings every 2 hours. Data from loggers provides a much more detailed record of groundwater fluctuations than the previous method of collecting data just once a month. Most importantly, the duration of water table depths at critical levels can be determined and correlations to surface flows can be derived.

Photo Stations

Ten photo stations were established throughout the study area with permanent numbered t-posts (Figure 3; waypoint locations are listed in Appendix A). Digital photographs were taken between mid-August and mid-September in 2003 through 2016 to visually document vegetation height, density, species composition, and overall site development. Annual photos were compared to evaluate visible changes over time.

Results

Avian Monitoring

Point Counts

Cleared/Overbank Area

Table C-1 in Appendix C provides data on the relative abundance of individual bird species for the Cleared/Overbank Area by year. The % *Plots* column shows the percentage of points in which the species was documented within this area. The *Mean* and *SD* columns represent the mean number and standard deviation of detections per point for the species.

There were 66 breeding bird species and 16 migrant species detected in the Cleared/Overbank Area during the point counts conducted from 2003 to 2016. During the first few years of monitoring, common breeding species (based on abundance and detection frequency) were red-winged blackbirds, blue grosbeaks, killdeer, western kingbirds, and brown-headed cowbirds. Common species by 2016 were yellow-breasted chats, spotted towhees, black-headed grosbeaks, mourning doves, blue grosbeaks, and black-chinned hummingbirds. These results are illustrated in the shade plot in Figure 7, which shows the average number of birds detected per point (relative abundance) of the most abundant species (generally ≥ 8 detections over the course of monitoring). The darker shades in each cell represent higher abundance at that sample point.

Species composition was analyzed using a Bray-Curtis similarity matrix which examines species similarity between years. Statistical analysis found a significant difference in species composition over time ($R=0.554$, $P<0.001$) within the Cleared/Overbank Area. Pairwise testing identified the highest similarities between years 2003 and 2004 and between years 2012, 2013, and 2014. For the most part, these results are illustrated in the Multi-dimensional Scaling (MDS) configuration in Figure 8 (note that the configuration may not exactly represent statistical results because MDS analysis uses means, unlike pairwise testing, and therefore variances may differ). MDS ordination ranks similarities and the associated configuration can be interpreted in terms of relative similarity of samples to each other (Clarke et al. 2014). For example, in this case it can be interpreted that species composition in 2005 and 2006 was less similar than all other years of monitoring. Species composition followed a continual change over time and began to become more similar starting in 2010 or 2011. Stress is the measure of distortion in the configuration. A stress factor of <0.5 gives an excellent representation; MDS analysis of this data had a stress of 0.07. The length and change in direction of the line between years illustrates the degree and relative change in species composition each year (e.g., starting in 2003 and ending in 2016). Size of overlay circles associated with each year represent abundance of 4 species, each of which was a species detected in the 4 most common




Figure 7. Shade plot of the most abundant species detected in the Cleared/Overbank Area by sample and year; darker shades in each cell represent higher abundance of that species. See Appendix B for species codes.




Figure 8. MDS ordination of 14 years of species abundance data based on Bray-Curtis similarities within the Cleared/Overbank Area (stress=0.07). Overlay circles associated with each year represent relative abundance of 4 of the species detected (BCHU=Black-chinned hummingbird, MODO=Mourning dove, YBCH=Yellow-breasted chat, RWBL=Red-winged blackbird).

guilds. In this case, abundance of black-chinned humming birds (edge guild) increased with time while abundance of red-winged blackbirds (water guild) decreased with time after peaking in 2005.

Table D-1 in Appendix D provides means and totals by species guilds for the Cleared/Overbank Area. Totals for the numbers of species within each guild accounted for all species detected during all three point count periods per year. Totals for the number of birds within each guild were calculated by averaging the number of birds detected at each point over the three point count periods and then summing all point averages. Note that sample sizes were sometimes different, so totals are not always equally comparable between areas or years. *Mean* and *SD* are the mean number and standard deviation of detections per point within each species guild.

The mean number of birds per point represents relative abundance (Nur et al. 1999), which is graphed by species guild over time in Figure 9. The total number of species detected during point counts represents species richness, graphed by guild over time in Figure 10. Since 2010, the most common species guilds based on relative abundance were midstory, ground shrub, and edge birds (Figure 9). There was an increase in both




Figure 9. Relative abundance by species guilds in the Cleared/Overbank Area over time. The number of total birds/point (red line) is graphed on the axis to the right.




Figure 10. Species richness by species guilds in the Cleared/Overbank Area over time. The total number of species (red line) is graphed on the axis to the right.

relative abundance and species richness among total birds over the monitoring period. Both of these variables increased in 2011 after a downward trend since around 2007. As of 2016, both relative abundance and species richness remained above 2011 levels.

Statistical analysis identified a significant relationship between relative abundance of birds (average number of birds per point) and time (year), in the total, cavity, mid-story, and water bird guilds (Table 1). In the total, cavity, and mid-story guilds there was an increasing trend in the relative abundance of birds detected; among water birds there was a decreasing trend. Although the P-value identified a difference in abundance over time for these bird guilds, low correlation coefficients (R values) indicated relatively weak relationships for all but the mid-story bird guild (see linear trend in Figure 11). An R value of 0.7437 indicated a moderately strong relationship between year and relative abundance among mid-story birds.

Table 1. P and R values for simple linear regression analysis between year and relative abundance by guild in the Cleared/Overbank Area. Alpha = 0.05.

Cleared/Overbank area 2003 to 2016		
Guilds	P-value	Correlation Coefficient [R]
Total birds	>0.001	0.2911
Canopy birds	0.100	0.1339
Cavity birds	>0.001	0.3424
Dense shrub birds	0.072	0.1465
Edge birds	0.078	0.1433
Ground shrub birds	0.530	-0.0514
Mid-story birds	>0.001	0.7437
Open birds	0.156	0.1157
Water birds	>0.001	-0.4933

Highlight = significant difference at the 95-percent confidence level




Figure 11. Linear trend in average number of mid-story birds per point in relation to year (2003 to 2016) in the Cleared/Overbank Area. Points represent the average number of observations within 3 repetitions at each point in each year, straight blue line represents best-fitting trend, and red curving lines represent 95 percent confidence intervals.

Burned Area

Table C-2 (Appendix C) shows relative abundance of individual species for the Burned Area by year. A total of 62 breeding bird species and 10 migrant species were detected in this area in 2003, 2004, and 2007 through 2016. The most common species detected in 2003 and 2004 (based on abundance and detection frequency) were turkey vultures, black-chinned hummingbirds, mourning doves, brown-headed cowbirds, spotted towhees, and yellow-breasted chats. By 2016 the most common species included black-chinned hummingbirds, yellow-breasted chats, spotted towhees, gray catbirds, mourning doves, and black-headed grosbeaks. The shade plot in Figure 12 shows the average number of birds detected per point (relative abundance) of the most abundant species over the course of monitoring. The darker shades in each cell represent higher abundance at that sample point. The pattern in species detections in the Burned Area appears to be more consistent over time than in the Cleared/Overbank Area (Figure 7) where there are varied breaks in species' detections over time.

Statistical analysis found a significant difference in species composition over time ($R=0.226$, $P<0.001$) within the Burned Area. Pairwise testing identified the highest species similarities between years 2003 and 2008; 2009, 2010, and 2012; 2011 and 2012; 2012 and 2013; and 2013 and 2014. These results are generally illustrated in the MDS configuration in Figure 13. The line between years illustrates relative change in species composition each year starting in 2003 and ending in 2016 with no data for years 2005 and 2006. In the Burned Area, MDS ordination shows species composition somewhat different in 2003, 2008, 2015, and 2016 from other years. This configuration had a stress of 0.11, which indicates an excellent representation. Size of overlay circles associated with each year represent abundance of 4 species, each of which was a species detected in the 4 most common guilds. It appeared that there was quite a bit of variability in the abundance of the 4 species throughout the monitoring period with no clear pattern of an increase or decrease in abundance. Species similarity analysis was the same done for Cleared/Overbank species composition, which is described in more detail above.

Means and totals by species guilds for the Burned Area are shown in Table D-2 (Appendix D). Relative abundance and species richness are graphed in Figures 14 and 15, respectively. Although there was variation in relative abundance between years, there were no major changes in the average number of birds detected from 2003 to 2016 (Table D-2, Figure 14). Species richness increased from 27 to 38 bird species detected over the study period.




Figure 12. Shade plot of the most abundant species detected in the Burned Area by sample and year; darker shades in each cell represent higher abundance of that species. See Appendix B for species codes.




Figure 13. MDS ordination of 11 years of species abundance data based on Bray-Curtis similarities within the Burned Area (stress=0.08). Overlay circles associated with each year represent relative abundance of 4 of the species detected (ATFL=Ash-throated flycatcher, BCHU=Black-chinned hummingbird, MODO=Mourning dove, YBCH=Yellow-breasted chat).




Figure 14. Relative abundance by species guilds in the Burned Area over time. The number of total birds/point (red line) is graphed on the axis to the right.




Figure 15. Species richness by species guilds in the Burned Area over time. The total number of species (red line) is graphed on the axis to the right.

In simple linear regression of abundance in relation to year only the open bird guild showed a significant trend in the relative abundance of birds detected, with the number of birds decreasing over time (Table 2). However, a relatively low R value indicated weak relationships between abundance and year for this species guild.

Table 2. P and R values for simple linear regression analysis between year and relative abundance by guild in the Burned Area. Alpha = 0.05.

Guilds	Burned area 2003, 2004, 2007 - 2016	
	P-value	Correlation Coefficient [R]
Total birds	0.430	-0.0642
Canopy birds	0.090	-0.1376
Cavity birds	0.841	-0.0163
Dense shrub birds	0.283	0.0873
Edge birds	0.098	0.1342
Ground shrub birds	0.838	0.0166
Mid-story birds	0.160	-0.1143
Open birds	0.032	-0.1731
Water birds	0.758	-0.0251

Highlight = significant difference at the 95-percent confidence level

Comparisons between Monitoring Areas

MDS ordination of species similarity including both monitoring areas is shown in Figure 16 (stress = 0.1). This perspective demonstrates that relative to the Cleared/Overbank Area, the Burned Area did not undergo large changes in species composition. The first




Figure 16. MDS ordination based on 14 years of square root transformed species abundance data and Bray-Curtis similarities for both the Cleared/Overbank and Burned Areas.

years of monitoring the two areas had very different species and with time, the Cleared/Overbank Area approached the Burned Area in species similarity. A statistical comparison determined there was a significant difference when comparing all years across both plots ($R=0.446$, $P=0.001$). Upon closer examination using pairwise testing between individual years and including both plots, no significant differences between 2012 and 2013 and between 2013 and 2014 were found.

Relative abundance was statistically compared between the two monitoring areas by years in which they were both sampled and by species guilds (see Table 3 for statistical results and P-values). In 2005 and 2006, the Cleared/ Overbank Area was the only site in which point counts were performed, therefore no comparisons between plots were made. Statistical comparisons between areas over time show that the Burned Area generally had a significantly greater number of total birds until 2012, when the Cleared/Overbank Area surpassed the Burned Area in relative abundance of total birds. In 2013 and 2014, total bird abundance in the two areas was statistically equal and in 2015 abundance in the Cleared/Overbank Area was again significantly greater than the Burned Area. In the early years of monitoring, the Burned Area usually had higher abundance of cavity, edge, and mid-story birds while the Cleared/Overbank Areas had higher abundance of dense shrub and water birds. By 2011, all guilds became statistically equal between areas with the exception of edge birds, which still had higher abundance in the Burned Area. There were no differences between areas in the abundance of birds in any guild in 2014 and 2016; in 2015 relative abundance of total, dense shrub, and water bird guilds was significantly

Table 3. Statistical comparisons of relative abundance between areas by year and guild. Alpha =0.05.

Year	Guilds								
	Total birds	Canopy birds	Cavity birds	Dense shrub birds	Edge birds	Ground shrub birds	Mid-story birds	Opening birds	Water birds
2003	P<0.001 ¹ Cleared<Burned	P=0.275 ²	P=0.006 ² Cleared<Burned	No dense shrub spp. in Cleared	P=0.329 ²	P=0.057 ¹	P<0.001 ² Cleared<Burned	P=0.578 ²	P<0.001 ² Cleared>Burned
2004	P=0.004 ¹ Cleared<Burned	No canopy spp. in Cleared	P=0.045 ² Cleared<Burned	P=0.938 ²	P=0.346 ¹	P=0.660 ¹	P<0.001 ² Cleared<Burned	P=0.059 ²	P<0.001 ² Cleared>Burned
2007	P=0.032 ² Cleared<Burned	No canopy spp. in Cleared	P=0.002 ² Cleared<Burned	P=0.005 ² Cleared>Burned	P=0.016 ¹ Cleared<Burned	P=1.00 ²	P<0.001 ¹ Cleared<Burned	No opening spp. in Burned plot	P=0.006 ² Cleared>Burned
2008	P=0.953 ²	No canopy spp. in Cleared	P=1.00 ²	P=0.015 ¹ Cleared>Burned	P=0.879 ¹	P=0.119 ¹	P=0.019 ¹ Cleared<Burned	No opening spp. in any plot	P<0.001 ² Cleared>Burned
2009	P=0.001 ² Cleared<Burned	No canopy spp. in Cleared	No cavity spp. in Cleared	P<0.001 ² Cleared>Burned	P<0.001 ¹ Cleared< Burned	P=0.704 ¹	P<0.001 ¹ Cleared<Burned	No opening spp. in Burned plot	P=0.004 ² Cleared>Burned
2010	P=0.033 ¹ Cleared<Burned	No canopy spp. in any plot	P=0.105 ²	P=0.010 ² Cleared>Burned	P=0.003 ² Cleared<Burned	P=0.309 ¹	P=0.130 ¹	No opening spp. in any plot	P=0.328 ²
2011	P=0.069 ¹	P=0.596 ²	P=0.668 ²	P=0.016 ¹ Cleared>Burned	P=0.017 ¹ Cleared<Burned	P=0.117 ¹	P=0.098 ¹	No opening spp. in any plot	P=0.200 ²
2012	P=0.032 ¹ Cleared>Burned	P=0.031 ² Cleared<Burned	P=0.063 ¹	P=0.006 ² Cleared>Burned	P=0.090 ¹	P=0.290 ¹	P=0.007 ¹ Cleared>Burned	No opening spp. in any plot	P=0.801 ²
2013	P=0.601 ¹	P=0.313 ²	P=0.133 ²	No dense shrub spp. in Burned	P=0.024 ¹ Cleared<Burned	P=0.067 ¹	P=0.293 ¹	No opening spp. in any plot	P=0.614 ²
2014	P=0.966 ¹	P=0.493 ¹	P=0.672 ¹	P=0.901 ²	P=0.920 ¹	P=0.929 ¹	P=0.170 ²	No opening spp. in any plot	P=0.569 ²
2015	P = 0.006 ² Cleared>Burned	P = 0.834 ¹	P=0.170 ¹	P = 0.030 ¹ Cleared>Burned	P=0.218 ¹	P=0.997 ¹	P=0.367 ²	No opening spp. in Burned plot	P=0.007 ² Cleared>Burned
2016	P = 0.609 ¹	P = 0.828 ¹	P=0.238 ¹	P=0.349 ¹	P=0.411 ¹	P=0.992 ¹	P=0.847 ¹	No opening spp. in Burned plot	P=0.525 ²

1=Student's t-test; 2=Mann-Whitney test of medians

Highlighted boxes = significant difference at the 95-percent confidence level

higher in the Cleared/Overbank Area.

Comparisons of trendlines and R^2 values for relative abundance and species richness between both monitoring sites are shown in Figures 17 and 18, respectively. Note that the R^2 values listed here were based on one number – the average number of birds or species detected per year – unlike those analyzed within each area and each guild, in which data from all points were used. Therefore, R^2 values differ. The Cleared/Overbank Area showed an increasing trendline for relative abundance (an increase that was determined to be statistically significant at $P < 0.001$) and species richness over time while the Burned Area showed almost no slope (i.e., no trend).

As can be seen on the graphs, in terms of actual values the Burned Area had consistently higher numbers of birds than the Cleared/Overbank Area. For example, in 2003 the relative abundance of total birds was 8.45 in the Burned Area compared to 2.75 in the Cleared/Overbank Area. This trend continued through 2011; in 2014 the Cleared/Overbank Area was equal to the Burned Area with both areas having an average relative abundance of 8.31. By 2015 the number of birds detected in the Cleared/Overbank Area (7.36) was higher than in the Burned Area (6.06; Tables D-1 and D-2). In 2016, relative abundance of total birds was essentially equal between the two areas (7.58 in Burned and 7.25 in Cleared/Overbank).

Willow Flycatcher and Yellow-billed Cuckoo Surveys

Willow flycatcher survey forms and maps are shown in Appendix E. In 2016, no SWFLs were detected within the boundaries of the LLRS. There was a total of 8 migrant WIFLs detected at areas adjacent to the LLRS between the Los Lunas and Belen bridges in Survey Site BL-25 (Figure 19) within Reclamation's Belen survey reach. Figure 19 also shows SWFL habitat suitability based on a model created for the Middle Rio Grande using 2016 vegetation maps (Reclamation, unpub. data). Most of the area between bridges is categorized as *Unsuitable* SWFL habitat with patches of *Moderately Suitable* habitat. The majority of the LLRS project area is classified as moderately suitable but the northwest corner (within the Burned Area) is considered unsuitable.

Yellow-billed cuckoo survey forms are also shown in Appendix E. There were no detections in either the BL-25 Survey Site or the LLRS in 2016. Since formal surveys began in 2014, one YBCU was detected within the LLRS project site in both 2014 and 2015 and two YBCUs were detected within the survey site in each of those years.




Figure 17. Trendlines and R^2 values for relative abundance over time in the Cleared/Overbank Area (2003-2015) and Burned Area (2003, 2004, 2007-2016).




Figure 18. Trendlines and R^2 values for species richness over time in the Cleared/Overbank Area (2003-2015) and Burned Area (2003, 2004, 2007-2016).




Figure 19. WFL detections and habitat suitability in the vicinity of LLRS within the Belen survey site (NAIP 2014 natural color photography).

Vegetation Monitoring

Vegetation Transects

Of the two areas included in avian point count monitoring, the Cleared/Overbank Area was the only area where vegetation monitoring was conducted throughout the entire study. As such, no comparisons were made between areas; only between years. In 2005 and 2006, survivorship of mixed shrub and cottonwood pole plantings was monitored at locations throughout the LLRS. Monitoring of mixed shrub and cottonwood pole plantings was discontinued once mortality/survivorship was documented. Fifty-four percent of the 160 mixed shrubs originally counted in 2005 at this site had survived by 2006 (Siegle 2007). New Mexico olive and Goodding's willow were the most successful species among the transplanted shrubs. The vast majority of cottonwood poles located within monitoring plots died (72 percent mortality). Based on recent observation, enough cottonwood poles were planted to result in long-term success of some trees but most cottonwoods onsite are due to natural regeneration.

Eighty annual and perennial plant species were detected in under- and overstory measurements during 14 years of vegetation monitoring. Common and scientific names of these species are listed in Table F-1 in Appendix F. Species richness at the site increased from 18 species detected in 2003 to 39 in 2016 and peaked at 44 in 2010 (Figure 20).




Figure 20. Plant species richness from 2003 to 2016.

Total percent cover by individual species, life-form (i.e., native or introduced shrubs < 1m, grasses, and forbs) and cover type (i.e., plants, litter, bare ground) of those species found in the understory layer are shown in Table F-2 in Appendix F.

Average total plant cover in the understory layer was variable over the course of monitoring, reaching a high of 79.6 percent in 2008 (Table F-2, Appendix F and Figure 21). From 2011 to 2016, total plant cover significantly decreased to levels comparable to those observed when monitoring began in 2003, which resulted in no change over the course of monitoring from 2003 to 2016 (Figure 21). Total cover of plant litter was 4.4 percent in 2003 and remained relatively stable until 2007. Since 2008 litter cover has




Figure 21. Statistical results analyzing total vegetation cover over time for various parameters. Red points represent mean, blue bars represent least significant difference intervals. Bars with the same letter indicate no significant difference while those with dissimilar letters indicate a significant difference in total cover between years ($\alpha=0.05$).

generally increased and peaked at 67.8 percent cover in 2012. Total litter cover was significantly less in the early years than in the later years (approximately 2009 to 2016) of monitoring. Total cover of bare ground decreased significantly over the monitoring period, from 63.5 percent in 2003 to 7.0 percent in 2016; bare ground was significantly higher in 2003 than in all other years (Table F-2, Appendix F and Figure 21).

Relative plant cover by life-form in the understory from 2003 to 2016 is shown in Figure 22. Native and introduced forbs and native grasses have been the predominant life-forms throughout monitoring with some shift in proportions from year to year.




Figure 22. Relative percent cover of life-forms in the understory layer from 2003 to 2016.

Understory shrub cover in 2006 (the year before measuring overstory as a separate layer) was higher than other years (Figure 21 and Table F-2). Shrubs over 1 m tall were still recorded in the understory yet this was the point that shrubs began reaching greater heights. All size classes of shrubs were included in 2006, which most likely led to higher values for understory shrubs than was truly representative. The regeneration of woody species, as represented by shrub cover in the understory layer, has remained stable over time with coyote willow and saltcedar typically the most common shrub species detected (Table F-2). In 2015 and 2016, a number of Siberian elm saplings were observed throughout the project area; the species made up 0.2 percent of the understory composition in transects. Native and introduced shrub species were relatively close in cover values, with native species generally having slightly higher cover in the understory layer. Native woody species (particularly coyote willow and cottonwood) have been more successful in maturing to the overstory layer.

Native grasses have sustained as a relatively high proportion of the understory composition throughout monitoring (Figure 22) and have apparently been successful in outcompeting introduced grasses at the LLRS. Native forbs have also sustained as a dominant lifeform at the site. Introduced forb cover was particularly high immediately after restoration activities and has remained one of the principal life-forms.

Total percent cover and average height of overstory species (woody species > 1 m in height) are shown in Table 4. Rio Grande cottonwood has continued to be the dominant woody species in the overstory canopy followed by coyote willow. Height estimates were gathered by measuring the tallest plants within the continual stretch of a species, therefore do not represent average heights of the stand but provide a consistent comparison from year to year.

The total cover of native overstory species significantly increased over time, expanding from 22.7 percent in 2007 to 96.3 percent in 2016, despite a significant drop from 80.9 percent in 2011 to 60.3 percent in 2012 (Table 4 and Figure 21). Total cover of

Table 4. Total percent cover and average height of woody overstory species (>1 m) from 2007 to 2016.

	2007		2008		2009		2010		2011	
Overstory plant species	Tot % cover	Avg ht (m)	Tot % cover	Avg ht (m)	Tot % cover	Avg ht (m)	Tot % cover	Avg ht (m)	Tot % cover	Avg ht (m)
Coyote willow	7.4	1.6	23.9	2.1	35.8	2.4	25.4	2.3	25.7	2.2
Goodding willow	0.3	1.6	0.9	2.4	1.5	2.9	1.0	3.3	1.0	3.4
Rio Grande Cottonwood	15.0	2.3	27.7	3.1	43.4	4.6	41.5	4.9	53.9	5.1
Narrowleaf cottonwood	0.0		0.0		0.0		0.0		0.3	5.3
Total native woody spp	22.7		52.5		80.7		67.9		80.9	
Saltcedar	4.3	2.3	5.8	2.2	9.7	2.8	8.9	2.8	6.5	2.6
Russian olive	0.6	2.9	1.1	3.4	1.6	3.9	1.9	5.2	2.5	4.7
Siberian elm	0.0		0.0		0.0		0.0		0.0	
Total introduced woody spp	4.9		6.9		11.3		10.8		9.0	
Total transect cover (accounting for overlap)	25.9		51.1		70.0		62.7		68.3	
	2012		2013		2014		2015		2016	
Overstory plant species	Tot % cover	Avg ht (m)	Tot % cover	Avg ht (m)	Tot % cover	Avg ht (m)	Tot % cover	Avg ht (m)	Tot % cover	Avg ht (m)
Coyote willow	14.2	2.3	22.2	2.4	23.1	2.4	32.0	2.5	30.2	2.3
Goodding willow	0.2	2.4	0.5	2.7	1.1	3.0	1.4	4.2	0.6	3.6
Rio Grande Cottonwood	45.4	6.4	49.9	6.4	53.8	7.1	55.1	7.8	64.8	8.9
Narrowleaf cottonwood	0.3	3.3	0.0		0.0		0.0		0.0	
Seep willow	0.2	1.9	0.0		0.0		0.1	1.8	0.7	2.3
Virgin's bower (vine)	0.0		0.0		0.2	2.5	0.0		0.0	
Total native woody spp	60.3		72.6		78.2		88.6		96.3	
Saltcedar	5.7	2.7	9.2	3.3	9.6	3.0	6.8	3.0	8.6	3.2
Russian olive	3.5	4.9	5.5	4.8	9.1	4.9	13.6	5.5	14.7	6.5
Siberian elm	0.2	2.8	0.2	2.9	0.3	3.7	0.1	1.6	1.2	2.9
Total introduced woody spp	9.4		14.9		19.0		20.5		24.5	
Total transect cover (accounting for overlap)	60.8		72.6		76.5		84.1		88.6	

introduced woody species was significantly greater in 2015 and 2016 than in 2007 and ranged from 4.9 percent to 24.5 percent. The overall transect canopy cover when accounting for overlap of species significantly increased from 2007 and 2008 to later years, peaking in 2016. Total canopy cover has followed a similar pattern to native overstory species since native species make up the majority of overstory canopy.

Since the onset of vegetation monitoring, the majority of plant species have been composed of native species relative to introduced in both the understory and overstory layers (Table 5). Relative cover of native understory species increased from 56 to 86 over the monitoring period while introduced species decreased from 44 to 14. Changes in relative cover of overstory species were not as drastic, with little variation over the monitoring period.

Table 5. Proportion of native and introduced species in the understory and overstory layers by year.

Year	Relative Percent Cover			
	Understory layer		Overstory layer	
	Native spp	Introduced spp	Native spp	Introduced spp
2003	56	44	NA	NA
2004	62	38	NA	NA
2005	74	26	NA	NA
2006	67	33	NA	NA
2007	74	26	83	17
2008	84	16	89	12
2009	89	11	88	12
2010	85	15	86	14
2011	76	24	90	10
2012	71	29	87	13
2013	75	25	83	17
2014	68	32	80	20
2015	83	17	81	19
2016	86	14	80	20

Analysis using a Bray-Curtis similarity matrix to compare plant species composition found a significant difference in species similarity between years ($R=0.624$, $P<0.001$). Pairwise testing identified the highest similarities between years 2011 through 2015. In general, these results are illustrated in the MDS configuration in Figure 23 (note that the configuration may not exactly represent statistical results because MDS analysis uses means, unlike pairwise testing, and therefore variances may differ). MDS ordination ranks similarities and the associated configuration can be interpreted in terms of relative similarity of samples to each other (Clarke and Warwick 2001). For example, in this case it can be interpreted that species composition in 2005 was less similar than that of all other years of monitoring. There was also a large difference in species composition from when monitoring began to the present. Stress is the measure of distortion in the configuration. A stress factor of <0.5 gives an excellent representation; MDS analysis of this data had a stress of 0.03. The line between years illustrates the degree and relative change in species composition each year (i.e., a very continual progression from 2003 to 2016 with species composition becoming more similar beginning around 2009). Size of overlay circles associated with each year represent average percent cover of the 3




Figure 23. MDS ordination of 13 years of plant species cover data based on Bray-Curtis similarities (stress=0.03). Overlay circles associated with each year represent percent cover of the 3 dominant overstory species (PODE=Rio Grande cottonwood, SAEX=coyote willow, TARA=saltcedar).

dominant overstory species each year. Total cover of the 3 species has increased with time, with larger increases in cottonwood and coyote willow.

Perennial pepperweed – a noxious weed – was documented at the site in 2003 and 2004, but inundation appeared to eradicate the species in 2005. In 2009, a patch of pepperweed was discovered between transect posts 3B and 4B and spotty occurrences of the weed were detected on the berm west of the river between transects 2 and 5. In 2010, perennial pepperweed total cover within transects peaked at 2.3 percent — up from minor detections in previous years (Table F-2). A patch was detected between transects 2 and 3 (about 1 acre) and pepperweed fell within transect 3. The patch between transects 3 and 4 had grown to approximately 2 acres in size. From 2011 to 2014, the cover of pepperweed within transects decreased, however occurrence of the species was noted in additional locations (between transects 1 and 2, on either side of mid-transect 6, and at transect 5). By 2015, the species had low occurrence on the berm. Perennial pepperweed appears to be confined to the north section of the site.

Vegetation Quantification Plots

Vegetation quantification plots were measured in August of 2016 within the Burned Area and in August of 2015 and 2016 within the Cleared/Overbank Area. When comparing data collected at LLRS sites, mean values within 0.5 standard deviations of mean values collected at nest sites were considered “suitable” for breeding SWFLs. For clarification, comparisons to all 112 nest sites will be referred to as “all” nest sites and comparisons to nest sites selected due to their similarity to the LLRS will be referred to as “selected” nest

sites. Of the 28 variables analyzed in this study, 13 were similar to *all* nest site values in the Burned Area and 10 were similar to *all* nest sites in the Cleared/Overbank Area in 2016 (values with * in Tables 6 and 7). These comparisons represent the best possible conditions for SWFL breeding habitat along the Middle Rio Grande. Ten variables were similar to *selected* nest sites in both the Burned and Cleared/ Overbank Areas (values in **bold** in Tables 6 and 7); these comparisons represent what are probably the most feasible conditions for the LLRS. Despite this assumption, the numbers of variables similar to comparison sites were about equal between *all* and *selected* sites, although “suitable” variables were not necessarily the same between comparison sites. The biggest differences in the two comparison populations were in species composition and in tree height and tree diameter at breast height in cm (DBH) Class II (higher in *all* nest sites), and shrub density and cover in the 0-3 m layer (higher in *selected* nest sites).

Table 6. Summary of center plot shrub and tree stem count data gathered at SWFL nest sites (2004 to 2006) and LLRS sites (2015 and 2016). Values in parentheses following nest means are “suitable” habitat ranges (+/- 0.5 sd). **Boldface values for LLRS sites are within “suitable” range compared to nest sites in selected reaches; values with * are within suitable range compared to all nest sites.**

Vegetation variable	All Reaches	Selected Reaches	LLRS Burned Area	LLRS Cleared/OB Area	
	Nest sites mean (n=112)	Nest sites mean (n = 22)	2016 mean (n = 3)	2015 mean (n = 3)	2016 mean (n = 3)
Shrub Stem Density (#/m ²)	3.64 (2.44 to 4.84)	5.62 (4.08 to 7.16)	2.40	4.56*	3.69*
Shrub Stem Spp Composition %					
<i>Salix gooddingii</i>	36.82 (17.52 to 56.12)	1.39 (0 to 3.85)	32.10*	0	0
<i>Salix exigua</i>	31.11 (13.81 to 48.41)	16.9 (3.40 to 30.41)	40.68*	56.54	66.94
Both <i>Salix</i> species	67.93 (49.23 to 86.63)	18.29 (4.99 to 31.59)	72.78*	56.54*	66.94*
<i>Populus deltoides</i>	1.26 (0 to 3.56)	2.28 (0.78 to 6.36)	0*	35.09	18.51
<i>Tamarix</i> sp.	23.15 (6.65 to 39.65)	50.24 (28.57 to 71.91)	16.70*	2.02	2.94
<i>Eleagnus angustifolia</i>	6.05 (0 to 15.6)	26.26 (11.02 to 41.51)	10.53*	6.35*	11.61*
Dead Shrubs %	37.00 (26.35 to 47.65)	33.10 (23.15 to 43.05)	65.97	29.19*	38.01*
Tree Stem Density (#/ha)	2,829 (2,164 to 3,494)	2,782 (1,979 to 3,586)	1,557	873	1128
Tree Stem Species Composition %					
<i>Salix gooddingii</i>	71.50 (52.35 to 90.65)	5.47 (0 to 12.30)	9.65	0	0
<i>Salix exigua</i>	5.09 (0 to 11.49)	0.78 (0 to 2.15)	0.60*	0*	0*
Both <i>Salix</i> species	76.59 (57.54 to 95.64)	6.25 (0 to 13.05)	10.24	0	0
<i>Populus deltoides</i>	3.36 (0 to 8.21)	7.42 (0 to 14.90)	1.02*	45.10	40.95
<i>Tamarix</i> sp.	11.93 (0 to 25.33)	49.14 (28.56 to 69.73)	54.20	0*	0*
<i>Eleagnus angustifolia</i>	8.12 (0 to 20.22)	37.20 (17.20 to 57.20)	34.53	54.90	59.05
Dead Trees %	3.96 (0.71 to 7.21)	7.31 (3.56 to 11.06)	13.60	9.32	11.11
Tree DBH Size Class Composition %					
Class 1 (5-10 cm)	70.06 (61.91 to 78.21)	78.71 (71.03 to 86.40)	67.84*	74.90*	55.93
Class 2 (10-20 cm)	29.02 (21.07 to 36.97)	18.91 (12.52 to 25.31)	28.95*	20.78	41.82
Class 3 (>20 cm)	0.92 (0 to 1.97)	2.38 (0.75 to 4.01)	3.22	4.31*	2.25

Table 7. Summary of point-centered quarter and canopy cover data from SWFL nest sites (2004 to 2006) and LLRS sites (2015 and 2016). Values in parentheses following nest means are “suitable” habitat ranges (± 0.5 sd). **Boldface values for LLRS sites are within “suitable” range compared to nest sites in selected reaches; values with * are within “suitable” range compared to all nest sites.**

Vegetation variable	All Reaches	Selected Reaches	LLRS Burned Area	LLRS Cleared/OB Area	
	Nest sites mean (n=112)	Nest sites mean (n = 22)	2016 mean (n = 3)	2015 mean (n = 3)	2016 mean (n = 3)
Shrub Canopy Layer					
Mean Plant Density #/ha	7,645 (3,776 to 11,515)	11,764 (6,083 to 17,424)	19,863	7,656	10,360*
Mean Plant Height	2.68 (2.28 to 3.08)	2.22 (1.54 to 2.90)	1.33	1.36	1.01
Mean Plant Crown Width	0.99 (0.82 to 1.17)	0.90 (0.51 to 1.29)	0.34	0.44	0.32
Canopy Layer					
Mean Plant Density #/ha	3,109 (1,941 to 4,277)	3,488 (1,912 to 5,064)	3,609*	5,311	11,671
Mean Plant Height	8.05 (7.27 to 8.84)	6.79 (6.22 to 7.37)	5.74	4.89	4.81
Mean Plant Crown Width	2.88 (2.36 to 3.40)	3.05 (2.36 to 3.74)	2.44*	1.40	1.50
Mean Cover Value					
0 – 3 m	28.70 (19.23 to 38.17)	37.51 (29.08 to 45.94)	40.25	49.25	31.75*
3 – 6 m	33.40 (23.77 to 43.03)	37.41 (28.65 to 46.18)	47.33	44.11	33.22*
>6 m	20.09 (11.49 to 28.70)	13.85 (8.91 to 18.79)	25.17*	18.19	15.58*

In reference to shrub and tree stem count data in Table 6, shrub stem density fell within the “suitable” range of *all* nest sites in the Cleared/Overbank Area. Shrub species composition was dominated by native willows (Goodding’s and coyote combined) in all samples except the *selected* nest sites, where willow was not a prevalent species. No Goodding’s willows were detected in the Cleared/Overbank Area at LLRS and in fact the percentage of coyote willow was above the suitability level when analyzed individually. In the Cleared/Overbank Area, combined willow composition fell within suitable range compared to *all* nest sites and percent composition of Russian olive was within suitable range of both nest site types. Cottonwood made up a much higher percentage and saltcedar made up a much lower percentage in the Cleared/Overbank Area than in nest comparison sites. Percent composition of all shrub species fell within the “suitable” range of *all* nest sites while no species fell within suitable ranges of *selected* sites in the Burned Area. The percentage of dead shrubs in the Cleared/Overbank Area was in the suitable range of both nest site types but was higher than suitable in the Burned Area.

Tree stem densities in both years were below the suitable range. No willow were recorded in the tree species composition in the Cleared/Overbank Area but the percentage of willow actually fell within the suitability range of *selected* nest sites (because selected sites had few willow in general) as well as in the suitability range of coyote willow composition of *all* nest sites. The percentage of cottonwood and Russian olive in tree species composition was above the suitable range of both nest site comparison samples. In the Burned Area, composition of all tree species fell within the suitable range of *selected* nest sites. The percentage of dead trees was higher than the nest site suitable ranges in both areas. With regards to tree DBH size classes, the percentage of trees with DBH > 20 cm fell into suitability range of *selected* nest sites in both areas; percent

composition of smaller DBH trees were suitable compared to *all* nest sites only in the Burned Area.

In reference to data collected using the point-centered quarter method in Table 7, plant density in the shrub layer was similar to both nest site samples in the Cleared/Overbank Area and shrub height and crown width was similar to both nest site samples in the Burned Area. Vegetative cover in the Cleared/Overbank Area met suitability standards of both *all* and *selected* nest sites at all height intervals and was similar to selected nest sites only in the 3 to 6 m interval in the Burned Area.

Groundwater Monitoring

Monthly Well Monitoring

Regular monthly well monitoring began in September 2004. The depth (in inches) below the ground surface to water at each well for each reading from June 2004 to October 2010 is summarized in Table G-1 in Appendix G. Data were used to create hydrographs that also included river discharge at the Rio Grande floodway in San Acacia, New Mexico (2003 to 2007) and at the Bosque Farms gauge (2008 to 2010; Figure H-1 in Appendix H). Discharge data collected near Los Lunas show flows in the Rio Grande are typically highest around April and May and lowest from July to September.

Within each transect (North, Middle, South as shown in Figure 3) groundwater levels varied. Water level within all wells was at ground surface level when discharges peaked around 4,600 cfs in May and June 2005. Wells along the South transect showed the largest differences in groundwater depth between wells compared to the Middle and North transects. The well nearest to the river (S1) was the shallowest and was rarely dry during monthly monitoring (Table G-1 in Appendix G). Groundwater at this well was less than 10 inches from the surface when discharges were greater than around 3,100 cfs and groundwater more than 50 inches from the surface when discharge fell below about 400 cfs. Well S2 (Figure 3) was typically dry at 61.5 inches during summer months (July-September) when river levels generally drop below 400 cfs.

The water table along the Middle transect was the shallowest measured, with Wells M1 – M3 rarely dry during monthly monitoring. The two wells nearest the river (M1 and M2) reached surface level when discharge was above approximately 3,200 cfs. The three wells nearest the river (M1-3) were relatively similar in groundwater depth, with groundwater at 15 inches or less from the surface when discharges were between 2,000 and 2,500 cfs. These wells only fell to more than 50 inches from the surface when the river was essentially dry.

Unlike the South and Middle transects, groundwater in the two wells nearest the river along the North transect where soils were sandy (N1 and N2) was generally deeper than in the two westernmost wells (N3 and N4). Clay soils at wells N3 and N4 most likely created shallow water table conditions and Well N3 was only dry in September 2003 and 2004 during monthly monitoring. When river discharge was between 3,200 and 3,500

cfs, groundwater depth was less than 10 inches from the surface in the shallower wells while the deeper wells were between 12 and 17 inches from the surface. The two shallower wells – N3 and N4 – only fell below 42 inches from the surface when the river was essentially dry.

Data Logger Well Monitoring

In June 2011, HOBO data loggers were installed. Groundwater data and river discharge at the gauge near Bosque Farms from June 2012 to September 2015 are graphed and included in Appendix H. Conditions were much dryer from 2011 to 2013 in the region, with peak flows only reaching about 1,700 cfs in April 2012. Flows rarely exceeded 750 cfs and the water table never reached the surface during this period.

Wells showed similar patterns in relative groundwater depth when comparing data from HOBO water level loggers with monthly data. Groundwater continued to be deepest at Wells S2, N1 and N2 with wells dry when river levels dropped below about 100 cfs. Well S2 was dry for most of the September 2012 to September 2013 period. All wells were dry from approximately August to November 2012. In July 2013, monsoons and associated increases in river discharge led to responses in groundwater level in all wells, though depths to groundwater and the length of time wells held water varied. Flows were much more consistent in 2014, with several peaks between 750 and 1000 cfs. The river was never dry and all wells held water throughout the year with the exception of Well S2 (groundwater present only when flows peaked) and Well N2 (rarely dry during summer months). A missing HOBO logger in Well N1 resulted in no data from September 2012 to September 2014. Flows were also fairly consistent in both 2015 and 2016 with peak flows much higher than in recent years – between 1,500 and 3,000 cfs. All wells held water during the monitoring period with the exception of Well S2, which was typically dry at 5.1 ft when flows fell below around 250 cfs. Wells M1, M3, N3, and S1 were less than 1.0 ft from the surface when flows peaked at 3,000 cfs in 2015; all of these loggers were stolen in 2016 and no data is available for this period. Loggers were refurbished and not operating from December 2014 through February 2015; therefore no data are available over this period. The Well M2 logger malfunctioned and no data were available for 2015.

The level of groundwater at the LLRS correlates closely with flows in the river, indicating a hydrologic regime influenced by the riverine system at the site (Appendices G and H). River discharge (which represents groundwater levels because the two are so closely linked) and vegetative cover are graphed in Figure 24.

Data loggers provided enough detail to discern diurnal fluctuations in the water table. Figure 25 shows an example of these fluctuations from September 2013 through August 2014. Groundwater fluctuated anywhere from 0.01 to 18.0 inches/day over a 3 year examination period (September 2011 through August 2014) at Well M2.




Figure 24. Hydrologic year (October – September) average discharge (cfs) in the Rio Grande at San Acacia (2002-2007) and at Bosque Farms (2008-2015), and the average total percent plant cover in transects at the LLRS, New Mexico. Restoration occurred in 2002; vegetation monitoring began in 2003. Overstory was not a separate cover measurement until 2007.




Figure 25. Diurnal fluctuation (ft) within Well M2 and average discharge (cfs) in the Rio Grande at Bosque Farms, New Mexico from September 2013 through August 2014.

Photo Stations

Photos taken from 2003 through 2016 are shown for comparison purposes in Appendix I.

Discussion

Avian Monitoring

Point Counts

Cleared/Overbank Area

Using the Burned Area for comparison, it appeared that desirable bird habitat developed over time within the Cleared/Overbank Area. By 2007, the Cleared/Overbank Area had higher numbers of dense shrub birds than the Burned Area, which was 5 years following restoration activities. In 2008 – 6 years after restoration – relative abundance became either statistically equal or greater than the Burned Area within all guilds except the mid-story guild, which had consistently greater abundance in the Burned Area. By 2010, relative abundance of mid-story species was equal between the two areas but total birds were greater in the Burned Area due to a significantly higher number of edge birds detected. From 2011– 9 years following restoration – to 2015, the two areas were essentially the same in relative abundance of most birds, although edge birds remained greater in the Burned Area. Species composition also became very similar between the two areas beginning in 2011 (Figure 18).

Increasing trends in relative abundance and species richness for cavity, dense shrub, and mid-story species guilds were consistent with the development of vegetation within the Cleared/Overbank Area, *i.e.*, as the cover and height of vegetation have increased (see Figure 26), so have the number and types of birds. Decreasing trends for opening and water birds are also consistent with habitat development patterns for these guilds; as the more open habitat required for these species has been replaced with denser vegetation, numbers of these birds have decreased.

Although most of the bird guilds in the Cleared/Overbank Area showed significant changes during the monitoring period, only the mid-story guild was found to show a strong statistically significant relationship with time with an R value of 0.7437, increasing from 2003 to 2016 (Table 1). The brown-headed cowbird was the most abundant species detected among mid-story birds until 2009, when the mean number of cowbirds detected per point dropped considerably. The brown-headed cowbird is not the most desirable of species because they use brood parasitism as a breeding strategy, which can reduce the productivity of host nests. Therefore, its decline may have been beneficial to other avian host species. Other mid-story species (e.g., black-headed grosbeaks, spotted towhees, and yellow-breasted chats) have increased, replacing the brown-headed cowbird as the dominant species in this guild. From 2003 to 2016, relative abundance of mid-story species increased from 0.17 to 4.10 birds/point and species richness increased from 3 to 12 (Table D-1), which are favorable trends for this site. The mid-story bird guild is an important indicator for the SWFL, which uses mid-story nesting habitat; therefore the increasing trend in mid-story species is an indication that the LLRS may be developing suitable habitat for SWFLs.




Figure 26. Development of vegetation at the LLRS as seen in 2002 immediately after the site was cleared (left) and in 2014 (right).

While the total number of birds in the Cleared/Overbank Area increased significantly from 2.75 birds/point in 2003 to 7.25 birds/point in 2016, only a weak linear relationship (R of 0.2911) was identified due to changing habitat and variable bird abundance. Relative abundance both increased and decreased over the study period as some habitat types declined while others became more developed. The number of total birds was closely linked to the number of water birds in this area until approximately 2009 (Figure 9). For example, the number of water birds peaked in 2005, when the LLRS was flooded, as did total number of birds. As vegetation in this area developed, habitat was less conducive to water birds. From 2009 to 2016, relative abundance of total birds closely correlated with the trend in mid-story birds (Figure 9). Further monitoring will determine if total birds continue to be more closely linked to mid-story birds with the development of this habitat type.

Burned Area

Results for the Burned Area were variable, indicating increasing and decreasing trends in both relative abundance and species richness among bird guilds, although none of the guilds showed exceptionally strong statistically significant relationships between abundance and year. This suggested that changes in bird populations may not have been strictly temporal and could have been caused by other factors affecting the site. A number of cottonwood snags have fallen since point counts were initiated, which changed the habitat somewhat and could be related to decreases in canopy birds. Relative abundance of mid-story birds was relatively high (4.69 birds/point) in 2003, three years after the fire. The average number of mid-story birds detected per point consistently decreased through 2010 (Table D-2 and Figure 14). Relative abundance has increased since then, ranging from 2.94 to 4.67 birds/point from 2011 to 2016. This was approximately the same period that mid-story bird abundance in the Cleared/Overbank Area began to steadily increase. The relatively high abundance of birds in this guild within both areas could be related to the development of habitat within the entire project area that is attracting more mid-story species in general.

Willow Flycatcher and Yellow-billed Cuckoo Surveys

It appears that suitable habitat has existed within adjacent sites between the Los Lunas and Belen bridges based on the occurrence of one SWFL territory in 2011, 2012, and 2013 (Moore and Ahlers 2015). Associated nests were successful in producing fledglings in 2012 and 2013. Much of the riparian habitat in the Belen survey reach is suitable as stopover habitat for migrating WIFLs as confirmed by presence/absence surveys; the number of resident SWFL territories detected within the reach has increased from 0 in 2009 to 20 in 2016. The 20 SWFL territories, which includes 13 breeding pairs, were found roughly 12 miles downstream of LLRS. This comprises the closest breeding population that could serve as a source for SWFL dispersal into the Los Lunas site.

Vegetation Monitoring

Vegetation Transects

A number of factors are important to the success of cottonwood/willow riparian forest restoration. These factors include soil conditions, such as salinity levels and texture, availability of native seed source, timing of high flows and flooding, and groundwater depth.

Alluvium texture is of primary importance in determining which plant species will succeed (Dressen et al. 2002). Lotic systems are characterized by fast moving water that deposits coarse alluvium of low fertility and high aeration. In contrast, lentic systems deposit fine alluvium (silts and clays) with higher fertility and less aeration. In general, lotic systems are conducive to the establishment of woody riparian trees and shrubs, while lentic systems are suitable for herbaceous wetland and marsh plants. The LLRS is a lotic system, as is the Middle Rio Grande bosque in general, although there are microsites where herbaceous wetland plants have established in depressions where silts and clays have deposited. In a restoration project on the Bosque del Apache National Wildlife Refuge (BDANWR), downstream of the LLRS, there was virtually no cottonwood germination in areas dominated by clay soils (> 65% clay), while regeneration of native species was greatest in sand deposits resulting from secondary channel development (Sprenger 1999) symptomatic of lotic systems.

Native species dominate the LLRS, particularly in the overstory, with cottonwood, coyote willow, and Goodding's willow present in the forest canopy. In the monitoring area, these species naturally re-established, indicating that a sufficient seed source was available on site. These species continue to regenerate, as is represented by shrub cover in the understory layer. Saltcedar and Russian olive are also re-establishing at the site. Saltcedar appears to be outcompeted by native willows and cottonwood which is a very positive outcome considering that saltcedar dominated the site when it was cleared, meaning there was an abundant seed source and resprouting potential for this species. The total percent cover of saltcedar after 14 years of monitoring was 0.8 percent in the understory (an indicator of the rate of regeneration) and 8.6 percent in the overstory, which is relatively low compared to other areas adjacent to the site. Evidence of *Diorhabda* spp. was detected in and around the LLRS in 2014 (Figure 27) and was apparent in photographs from photo stations 6 through 10 in 2016 (Appendix I). This beetle was released in 2001 at several sites across the Southwest as a biological control for saltcedar and is spreading into areas beyond its predicted range, including the Middle Rio Grande. The effects from *Diorhabda* could potentially reduce saltcedar, an outcome that monitoring would detect. Saltcedar that fell within the vegetation transects did not show signs of beetle forage from 2014 to 2016. Russian olive, another introduced species, has been gradually increasing in cover over time and now composes 14.7 percent of overstory cover.

Of course, although a local natural seed source is important to successful restoration, it must be combined with hydrologic conditions optimum for cottonwood and willow



Figure 27. Evidence of *Diorhabda*, a biological control beetle released in the Southwest to manage saltcedar, was first observed in the LLRS in 2014.

regeneration and establishment. A restoration site in the urban Albuquerque reach of the Middle Rio Grande used a design similar to the one implemented at LLRS by incorporating natural hydrologic processes; 10,000 cottonwoods/ha established at this site following overbank flooding as compared to a higher site out of reach of the flood in which no trees established following the same event (Muldavin et al. 2015). Not only is overbank flooding necessary, it must be timed with germination of willow and cottonwood seedlings. Investigations at the BDANWR proved that natural recruitment of willow and cottonwood was possible subsequent to over-bank flooding during peak river flows in late May and early June (Sprenger 1999). Flooding conditions at LLRS were apparently conducive to natural recruitment of native species, especially from 2005 to 2009 when average annual discharge rates were relatively high compared to other years (Figure 24). The rate of stream stage decline should not exceed 2.5 cm per day for seedling survival (USDA, NRCS 1998), a criterion that was presumably met. Cottonwood and willow seedlings were detected early in the study, starting in 2003 which was the first year of vegetation monitoring. Establishment of woody species, however, was especially evident during the 2006 growing season, the year after extremely high river flows and prolonged flooding on site. The length of inundation from flooding also affects the ability of plants to germinate and sustain. Mortality of cottonwoods submerged for over 32 days was 100 percent in studies by Sprenger (1999) and Hosner (1958 as cited by Sprenger 1999). Coyote willow, on the other hand, was found to survive after 2 months of inundation in New Mexico (USDA, NRCS 1998). Monthly groundwater well data collected in this study did not provide enough detail to determine how long flooded conditions persisted at the LLRS. From 2011 to 2014, when more complete groundwater data was collected with HOBO logger instruments, no

flooding occurred. Hydraulic modeling of the LLRS determined that discharge of 2,500 cfs (design goal) would cause extensive inundation of the site (Kissock 2010). The water table reached the surface for approximately a week in May 2015 and in June 2016 when flows peaked between 2,500 and 3,000 cfs (Appendix H).

Depth to groundwater plays a key role in determining which riparian species will succeed in a restored site. The primary rooting zone for obligate riparian plants is the capillary fringe above the water (Dressen et al. 2002). The thickness of the capillary fringe is controlled by soil texture, with finer textured alluvium having a broad zone of unsaturated soil with high moisture content. A thicker capillary fringe zone has a greater water content however it also has lower aeration resulting from less air-filled pores. Because woody riparian species generally require highly aerated soils, suitable restoration sites generally have a thin capillary fringe with lower water content but more air filled pores. Groundwater conditions at the LLRS are discussed in the Groundwater Monitoring section below.

Vegetation Quantification Plots

Some portions of the Cleared/Overbank and Burned Areas may have developed riparian vegetation of suitable height, density, and structure to provide breeding habitat for the SWFL. Based on both avian and vegetation monitoring, the area has been productive in terms of developing native overstory habitat, and SWFLs could potentially occupy the LLRS in time. Unfortunately, it is difficult to accurately assess the habitat suitability of a site for breeding SWFLs based solely on visual observations since the factors that appear to influence site selection are numerous and variable. Vegetation quantification data was collected within the LLRS in 2015 (Cleared/Overbank Area) and in 2016 (Cleared/Overbank Area and Burned Area) in an effort to evaluate habitat for SWFL breeding (Figure 4).

In 2007, sites in the Burned Area of LLRS were compared to similar data collected from sites downstream where SWFL nests were known to occur (Moore 2009). At that time, vegetation at the Los Lunas site was found to be more dense, and of a younger age-class than sites where SWFL breeding took place. It was determined that the Los Lunas site would more closely approximate occupied SWFL breeding habitat in “a few growing seasons.” Based on visual observation, small isolated patches of vegetation likely reached structural suitability around 2010 within the Burned Area.

In 2015, 3 plots in the Cleared/Overbank Area were sampled and comparisons were expanded to include not only the original 112 nest sites (ideal habitat) but also selected nest sites that may be a better representation of LLRS potential to develop into suitable habitat (data from this sampling period is included in Tables 6 and 7).

In 2016, the vegetation quantification study was augmented to include 3 plots in the Burned Area. Based on data from this year, the Cleared/Overbank Area did not provide optimal SWFL habitat, particularly in relation to characteristics found in the tree layer. Shrub density fell within suitability ranges of nest sites, although average shrub height and crown width were less than suitable. The shrub species composition of Goodding's

willow, combined willow and Russian olive all met suitability standards of at least one of the comparison nest sites. Closer examination of this data, however, reveals that the percent composition of coyote willow was higher than suitable, and because no Goodding's willow was detected in this area, the combined willow composition was strictly coyote willow, which was determined to make up a higher proportion of the composition than is desirable. Trees in the Cleared/ Overbank Area were too dense and too small to be considered suitable habitat in 2016. The only tree species recorded in this area were cottonwood and Russian olive; both comprised more of the species composition than was considered suitable. Even though willow and saltcedar were not present in the tree layer, a composition of 0 percent fell within suitability ranges of selected nest sites (Goodding's and coyote willow) and all nest sites (saltcedar). The percentage of trees with DBH > 20 fell into suitability range of selected nest sites; the percent composed by smaller DBH trees were either too low (0 to 10 cm) or too high (10-20 cm) to fall within suitability ranges. Percent cover within all height intervals was similar to both nest site comparison data. Although some conditions were met, species composition and tree density and height in the Cleared/Overbank Area were factors that appeared to be the most limiting. The high percentage of cottonwood at the LLRS may inhibit development of optimum SWFL habitat if the site matures into a cottonwood gallery. However, cottonwood could potentially contribute to desirable habitat for the YBCU. The amount of data collected was limited (n= 3) and a stronger analysis could be made with more samples.

Vegetation quantification data collected within the Burned Area also suggested that SWFL habitat was not necessarily ideal but was somewhat better than in the Cleared/ Overbank Area. Physical characteristics of the shrub layer did not meet suitability standards but shrub species composition was within the suitability range of all nest sites for all species. Tree density (point-centered quarter measurement) and crown width both fell into the suitability ranges of both comparison nest site types; tree height was considered to be too short. As with shrub species, all tree species were within the suitable range of percent composition (mostly selected nest sites). All classes of DBH were found to be similar to one of the two comparison nest site samples. Finally percent cover in the 0 to 3 m interval was similar to selected nest sites and the > 6 m interval was similar to all nest sites; percent cover within the 3 to 6 m range was higher than nest site samples. In the Burned Area, species composition in both the shrub and tree layers appears to be adequate, as do most of the tree characteristics (with the exception of a tall enough canopy and thicker vegetation from 3 to 6 m height). The shrub layer did not appear to be developed enough to provide optimal habitat as compared to occupied areas.

Habitat suitability modeling in 2016 determined the LLRS site to be mostly *Moderately Suitable* (approximately 77 acres) with some *Unsuitable* (approximately 42 acres within the Burned Area) based on Hink and Ohmart (1984) vegetation classification (Reclamation, unpub. data). The Cleared/Overbank Area was characterized as a cottonwood overstory (15-40 ft average) with a coyote willow or coyote willow-saltcedar understory. Vegetation types were more variable in the Burned Area, with cottonwood 15-40 ft over Russian olive –saltcedar in the northern portion, cottonwood greater than 40 ft over Goodding's willow-coyote willow and Russian olive-saltcedar 15-40 ft over

coyote willow in the southern portion. All vegetation types within the LLRS were estimated to have aerial cover greater than 50 percent. The limiting factor with the Hink and Ohmart classification is that the density and structure by layer is unknown based solely on the label. In this case, the vegetation quantification data provided detailed information. Both types of data appear to indicate that although conditions provide fairly good SWFL habitat there are still limitations.

Groundwater Monitoring

Groundwater depth at the LLRS correlated closely to Rio Grande flows (Appendices G and H), indicating that connectivity between the shallow aquifer and the river is still functioning despite management activities that could potentially impact hydrologic processes such as channelization, regulation of surface flow, groundwater pumping, and water diversions. Because flows influenced the water table depth, total percent plant cover also correlated with river discharge rates (Figure 24), particularly shallow-rooted understory plant species. There were shifts in understory vegetation composition (see 2005 and 2006 in Figure 22) as well as noticeable increases in growth in 2006 following the extended period of inundation in 2005. Flooded conditions led to germination and establishment of riparian plants (especially coyote willow and cottonwood as demonstrated in Table F-2, Appendix F). The relatively high discharge rates in 2008 did not lead to long periods of inundation, but did result in a high water table. These conditions provided plant available water and allowed for increased plant cover that year. Yearly discharge rates decreased after 2008 but have been steadily increasing since 2014; understory vegetative cover has followed a similar pattern.

Overstory cover remained somewhat stable from 2009 to 2013 despite decreasing discharge rates. This would suggest that by 2009, cottonwood and willow had developed a deep enough root system to sustain declines in the water table. Regardless, based on well monitoring data, it is unlikely that groundwater at the site has fallen below the crucial depth of around 10 ft necessary to sustain woody riparian species (Cartron et al. 2008). Most wells, which average around 5 ft in depth, were only occasionally dry, which indicates that the water table is relatively shallow at the site. On the other hand, vegetation did appear to be affected by prolonged dry conditions at the site. From 2010 to 2012, overstory foliage was observed to be rather sparse and leaves were dropping earlier than expected. This is supported by overstory cover values, which did not notably increase from 2009 to 2012. Since 2013 overstory cover has been gradually increasing along with increasing discharge rates.

The three wells nearest to the river and within (or near) the vegetation monitoring site show that groundwater is deeper in the northern section of the site. Groundwater depth did not appear to have a direct correlation with overstory vegetation cover, which was relatively consistent throughout transects. There was a small effect on species composition based on Hink and Ohmart vegetation types. Saltcedar was prevalent enough to be included in understory classification in the southern section where the water table is shallower. These results imply that although the water table falls below well depth more

frequently in the north, differences in groundwater depth are not great enough to result in large variability in vegetation.

Data from the HOBO water level loggers were collected every 2 hours from June 2011 to September 2016, which captured diurnal fluctuations in the water table (Figure 25). Diurnal fluctuation in shallow water tables is attributed to groundwater consumption by phreatophytes such as willow and cottonwood (Shah et al. 2007). The significant evapotranspiration (ET) consumption of phreatophytic plants influences the behavior of interconnected surface and groundwater systems. The water table, which declines rapidly during daylight due to ET, partially recovers at night. The recovery in the evening and night hours is attributed to lateral and vertical groundwater flow to the discharge area (Shah et al. 2007). As Figure 25 shows, in many cases a spike in river discharge also caused a spike in diurnal fluctuation, indicating that river flows were controlling fluctuations in well depth. In general, diurnal fluctuations were highest during the growing season (approximately May through September), which is a representation of plant ET at the site.

The ET of surrounding plant species influences diurnal fluctuations in association with groundwater depths. Transpiration by mature cottonwood is unaffected as long as the water table is within 3 m of the surface (Cleverly et al. 2006). When groundwater is drawn down deeper, transpiration declines with increasing crown dieback. Goodding's willow is found in habitats similar to those of cottonwood; therefore Goodding's willow ET is expected to respond to groundwater depth in the same manner as cottonwood (Cleverly et al. 2006). Conversely, coyote willow can tolerate dryer conditions, much like saltcedar, and ET from coyote willow is expected to respond more like saltcedar. Saltcedar transpiration is not restricted by depth to groundwater as it is in cottonwood (Cleverly et al. 2006). Even though saltcedar ET is not dependent upon depth to the water table, it does respond to changes in water table depth, increasing while the groundwater is falling.

Data collected at LLRS is not specific enough to correlate individual wells with surrounding plant species. Another limitation is that wells do not go to depths that are found to inhibit ET of cottonwood and Goodding's willow. Nonetheless, patterns in diurnal fluctuations are apparent. There is an increase in diurnal fluctuation while river levels fall, which could indicate coyote willow ET responding to a deepening water table, or it could simply be a seasonal pattern (i.e. summer months are the growing season and also when river flows decline).

Photo Stations

Photos taken at Stations 1 through 5, which are located along the berm and face east toward the river in the Cleared/Overbank Area where vegetation transects are located, show considerable and steady growth in regenerating willow and cottonwood. In 2006, following flooded conditions in 2005, the establishment of woody species appears stable. By 2009, a definite overstory has developed. From about 2010 through 2012, foliage is noticeably affected by dry conditions and there is not an obvious growth in overstory

species. In photos taken at Stations 6 – 10, which are located along the road and face east toward the Burned Area, the density of standing dead cottonwoods in the burned forest has noticeably decreased over the years as the growth of regenerating understory has increased. This is the area in which cottonwood poles were planted in 2004, and a healthy stand of cottonwoods is developing in this area. Saltcedar is also evident in many of the photos. In 2016 photos, foliar impacts to saltcedar from *Diorhabda* become distinct.

From 2010 to 2012, it was observed that leaves were already turning yellow and beginning to fall during monitoring in early to mid-September, which may have been due to an extended period of low precipitation (Figure 24). This condition is apparent in photos from these years. By 2013, despite continued drought, foliage is fuller and greener (which was supported by data that showed an increase in overstory cover this year). Tree leaf cover appeared sparse in 2015 (Figure 28 and Photo Stations 1-5, Appendix I), with leaves falling by early September; reasons are unknown but may be long-term effects of drought in the region. Data did not reflect this with overstory cover higher in 2015 (84.1 percent) than in previous years. By 2016 there were no observable declines in overall vegetation health at the site.



Figure 28. Example of sparse leaf cover on a cottonwood, September 2015, LLRS.

Conclusion and Recommendations

Avian Monitoring

Conclusions

Avian relative abundance and species richness data have been collected for a 14 year study period at the LLRS in riparian habitat along the Middle Rio Grande. Monitoring has tracked the development of the avian population and of SWFL habitat suitability in

the restoration area where established stands of native riparian vegetation bordering high flow channels is the desired future condition.

Despite decreasing trends in relative abundance of total birds in both avian monitoring areas from approximately 2005 to 2009, bird detections have either maintained (i.e., Burned Area) or increased (i.e., Cleared/Overbank Area) from 2003 to 2016. These results are similar for species richness in each monitoring area as well. The reasons for decreases mid-study are unknown, but regardless, riparian habitat in the LLRS currently appears to be supporting diverse avian populations.

The abundance and diversity of breeding cavity, dense shrub, ground shrub, and mid-story bird species in the Cleared/Overbank Area have increased during monitoring, resulting in an overall increase in total bird abundance. The mid-story guild serves as an indicator for SWFL habitat. Since 2010, the number of mid-story species detections per point in the Cleared/Overbank has been either statistically equal to or greater than the Burned Area. Both areas (restored and burned) appear to show promising potential for providing SWFL habitat. As woody riparian plants develop height and density suitable for nesting substrate and cover in the Cleared/Overbank and Burned Areas, mid-story habitat vital to SWFLs should continue to increase.

Based on avian data collected in this study, mid-story habitat – potentially suitable for SWFL breeding - became established by approximately 2010. Using the LLRS as a reference, it appears that it is possible for SWFL habitat to develop within 8 years following restoration activities in the Middle Rio Grande. These results are dependent on environmental conditions that are favorable for successful restoration, as were present during development of this site. Using hydraulic and geospatial analysis, Kissock (2010) determined that the LLRS is “sustainable by continuing to provide habitat to endangered species targeted for habitat restoration (i.e. SWFL and minnow).”

Based on vegetation data collected specific to SWFL habitat, by 2016 certain variables related to overstory species composition and structure were not comparable to occupied nesting sites but nonetheless many conditions had been met. Although samples were limited (n=3), this data does provide a general idea of limitations in SWFL habitat at LLRS. Habitat evaluations could be improved with more vegetation quantification data. Over the past several years, SWFLs have established territories in closer proximity to the LLRS, increasing the likelihood that they may occupy the site in the near future.

Recommendations

Continue avian monitoring in accordance with the initial monitoring requirements of the BO and to provide information for adaptive management of SWFL restoration projects. Further monitoring will help to determine if the Cleared/Overbank Area can sustain habitat for most bird guilds, especially for mid-story species that include the SWFL. It is also important to document occurrence of breeding SWFLs at the LLRS to determine if suitable habitat has in fact developed, which was one of the objectives for restoration of this site.

Vegetation Monitoring

Conclusions

Vegetation monitoring data are being used to document:

- 1) the natural establishment of riparian vegetation in the disturbed areas
- 2) the establishment of wetland vegetation in depression areas
- 3) the possible establishment of noxious weeds and recolonization of exotics, and
- 4) rates of vegetation development for future SWFL restoration efforts.

Success of riparian restoration at the LLRS could also potentially be used for comparison at other restoration sites along the Middle Rio Grande.

Riparian vegetation has successfully established in the Cleared/Overbank Area. Native species dominated the overstory and included coyote willow, Goodding's willow, and Rio Grande cottonwood. The wetland indicator status of both willow species is "facultative wetland" (i.e., usually occur in wetlands but may occur in nonwetlands) based on the National Wetland Plant List for the Arid West (USDA NRCS 2014). In the understory layer, native species also dominated the vegetation, although were not necessarily considered riparian plants. The native grass vine mesquite, for example, was the most common understory species detected at the site and is in the "upland" wetland indicator category. Plant species found in depressions, however, were categorized as "obligate wetland" (i.e. almost always occurs in wetlands, e.g., common spikerush) or as "facultative wetland" (e.g., fragrant flatsedge, Baltic rush, common reed, and sword-leaved rush). Saltcedar, although present at the site, had relatively low cover values (<10 percent) over the monitoring period and did not appear to be competitive with native overstory species.

Prichard et al. (1998 as cited in Dressen et al. 2002) developed a comprehensive assessment of criteria useful in judging riparian area condition and attributes that constitute a proper functioning condition for lotic areas. The vegetation attributes of a proper functioning riparian system include:

- 1) the age class distribution of the riparian plant community indicates the recruitment of young individuals and the maintenance of older individuals;
- 2) the species composition of the riparian area is diverse;
- 3) the characteristic soil moisture of a riparian-wetland area is indicated by the species present;
- 4) species with root masses capable of protecting against high flow events are present on the streambanks;
- 5) the condition of the riparian plant community is healthy and robust;
- 6) vegetative cover is sufficient to protect streambanks and dissipate energy during high flow events; and,
- 7) the riparian plant community can provide sufficient large woody debris to act as an agent to modify the hydrology if necessary for proper functioning.

When evaluating the LLRS using these attributes, most of these criteria appear to have been met. Tree and shrub species detected in the understory layer are an indication that woody species are regenerating at the site and have been throughout monitoring. A diverse composition of riparian species, including willow, cottonwood, sedges, and rushes, are present. The condition of vegetation appears healthy. Even during drought conditions, canopy cover maintained at a stable rate, which also indicates that woody vegetation has reached rooting depths that can sustain a deeper and fluctuating water table. Woody debris is present in the form of downed cottonwood as a result of the fire in 2000. High energy flows and prolonged inundation occurred in 2005 and flooding was again recorded for short periods in 2015 and 2016; the site appeared to withstand these events without major impacts.

Conditions that are important to the success of riparian restoration, which include groundwater depth, timing of high flows and flooding, native seed source, competition from exotics, and soil conditions (i.e., texture and salinity levels) have all been conducive to development of healthy, native riparian habitat. In conjunction with favorable conditions, the techniques used for restoring the site can also be deemed successful thus far. The success of restoration at this site can largely be attributed to a design that integrated natural hydrologic processes; banks were lowered to allow for overbank flooding and channels created to slow flood waters and encourage sediment deposition (Muldavin et al. 2015). Kisko (2010) predicted that the LLRS would require maintenance in the future due to greater than critical shear stress values, resulting in a tendency towards erosion. At this point in the study, erosion does not appear to be problematic.

Recommendations

Monitoring should be continued at the established vegetation transects in accordance with the initial monitoring requirements of the BO and to provide information for adaptive management of SWFL restoration projects. Long-term monitoring will help to determine if vegetation at the site can continue to regenerate and sustain varying conditions.

In an attempt to specifically evaluate the site for SWFL habitat suitability, continue nest site quantification studies in both the Burned and Cleared/Overbank Areas to increase sample size and more accurately estimate habitat conditions for the species.

As of 2010, large patches of perennial pepperweed were detected within the LLRS. The occurrence of this noxious weed has expanded from previous years. Also, based on general observation and supported by cover data, Russian olive has noticeably increased throughout the area. A number of Siberian elm seedlings and saplings were also observed in 2015 and 2016. Control of these species may be warranted.

Groundwater Monitoring

Conclusions

Data from monitoring wells were used to correlate the development and extent of wetland/riparian type vegetation at the restoration site. These data have been instrumental in interpreting long-term development of plant communities at the LLRS. The depth of the water table has a large effect on the continued success of cottonwood and willow. For example, Hultine et al. (2010) found that cottonwood has a greater sensitivity to interannual reductions in water availability, while willow is more sensitive to longer periods of soil water depletion.

It appears that the water table at the LLRS is relatively shallow, which has been important in recruiting and establishing stands of cottonwood and willow. Most of the wells, all of which averaged around 5 ft in depth, held water throughout the majority of the year. Based on groundwater data and on the development of healthy native vegetation, it is unlikely that the water table falls to depths that are detrimental to the success of woody riparian species. Vegetation did, however, appear to show stress from dry conditions in recent years.

Recommendations

Data from water level loggers is useful in determining groundwater effects on developing vegetation and associated wildlife habitat at the site, as well as evaluating the connectivity of groundwater and surface water flows. Groundwater monitoring should be continued for the duration of the study, particularly in light of dry conditions that have occurred in recent years.

Photo Stations

Conclusions

Shifts in plant composition and growth stages of regenerating willow and cottonwood have been observed over the 14 years of monitoring. Photos have provided an important record of the changing vegetation, including the timing of certain stages in development. Of all the methods of data collection used, photographic documentation has probably presented the clearest account of the changes at the LLRS.

Recommendations

Trends in the vegetation should continue to be captured through photos for the duration of the study.

Literature Cited

- Ahlers, D. and L.H. White. 1997. Southwestern Willow Flycatcher study results – 1996: selected sites along the Rio Grande from Velarde, New Mexico, to the headwaters of Elephant Butte Reservoir. Bureau of Reclamation, Technical Service Center, Ecological Planning and Assessment. Denver, CO.
- Army Corps of Engineers. 2000. Installing Monitoring Wells/Piezometers in Wetlands. Wetlands Regulatory Assistance Program. ERDC TN-WRAP-00-02. July 2000.
- Cartron, J.C., D.C. Lightfoot, J.E. Mygatt, S.L. Brantley, and T.K. Lowrey. 2008. A Field Guide to the Plants and Animals of the Middle Rio Grande Bosque. University of New Mexico Press. 375 pp.
- Clarke, K.R., R.N. Gorley, P.J. Somerfield, and R.M. Warwick. 2014. Change in Marine Communities: An Approach to Statistical Analysis and Interpretation, 3rd edition. Plymouth, United Kingdom.
- Cleverly, J. R., C. N. Dahm, J. E. Allred Coonrod . 2006. Groundwater, Vegetation, and Atmosphere: Comparative Riparian Evapotranspiration, Restoration, and Water Salvage. USDA Forest Service Proceedings RMRS-P-42CD.
- DeRagon, W., P. Mehlhop, A. Henry, and D. Mehlman. 1995. Habitat measurements of Southwestern Willow Flycatcher nest sites along the Rio Grande in New Mexico, 1995. U.S. Army Corps of Engineers. Albuquerque, NM.
- Dreesen, D., J. Harrington, T. Subirge, P. Stewart, and G. Fenchel. 2002. Riparian Restoration in the Southwest: Species Selection, Propagation, Planting Methods, and Case Studies. In Dumrose, R.K., Riley, L.E., and Landis, T. D., technical coordinators. National proceedings: forest and conservation nursery associations – 1999, 2000, and 2001. Proceedings RMRS-P-24. Ogden, UT: U.S. Department of Agriculture Forest Service, Rocky Mountain Research Station. P. 253-272.
- Halterman, M., M.J. Johnson, J.A. Holmes and S.A. Laymon. 2015. A Natural History Summary and Survey Protocol for the Western Distinct Population Segment of the Yellow-billed Cuckoo: U.S. Fish and Wildlife Techniques and Methods, 45 p.
- Hink, V. C., and R. D. Ohmart. 1984. Middle Rio Grande biological survey. Army Corps of Engineers Contract No. DACW47-81-C-0015. Albuquerque, NM
- Hultine, K. R., S. E. Bush, and J. R. Ehleringer. 2010. Ecophysiology of Riparian Cottonwood and Willow Before, During, and After Two Years of Soil Water Removal. Ecological Applications 20:347–361.

- Kissock, S. 2010. Hydraulic and Geospatial Analysis of Stream Engineering and Habitat Restoration near Los Lunas, NM. Master's Thesis in Civil Engineering, University of New Mexico.
- Martin, T.E., C.R. Paine, C.J.Conway, W.M. Hochachka. P. Allen, and W. Jenkins. 1997. BBIRD field Protocol. Montana Cooperative Wildlife Research Unit. University of Montana. Missoula, MT.
- Moore, D. 2009. An Assessment of Potential Southwestern Willow Flycatcher Habitat. Bureau of Reclamation, Denver, CO.
- Moore, D. 2007. Vegetation quantification of Southwestern Willow Flycatcher Nest Sites. Bureau of Reclamation, Technical Service Center, Fisheries and Wildlife Resources. Denver, CO.
- Moore, D., and D. Ahlers. 2016. 2015 Southwestern Willow Flycatcher Study Results: Selected Sites along the Rio Grande from Bandelier National Monument to Elephant Butte Reservoir, New Mexico. Bureau of Reclamation, Denver, CO.
- Muldavin, E., E. Milford, N. Umbreit, and Y. Chauvin. 2015. Long-term Outcomes of a Natural-processes Approach to Riparian Restoration in a large Regulated River: the Rio Grande Albuquerque Overbank Project After 16 Years. U.S. Department of the Interior, Bureau of Reclamation, Albuquerque, NM.
- Nur, N., S.L. Jones, and G.R. Geupel. 1999. A Statistical Guide to Data Analysis of Avian Monitoring Programs. U.S. Department of the Interior, Fish and Wildlife Service, BTP-R6001-1999, Washington, D.C.
- Poff, B., K.A. Koestner, D.G.Neary, and D. Merritt. 2012. Threats to Western United States Riparian Ecosystems: A Bibliography. Gen. Tech. Rep. RMRS-GTR-269. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 78 p.
- Shah, N., M. Nachabe, and M. Ross. 2007. Extinction Depth and Evapotranspiration from Ground Water under Selected Land Covers. Ground Water 45:329–338.
- Siegle, R. 2007. 2006 Monitoring Report for the Los Lunas Habitat Restoration Site. U.S. Department of the Interior, Bureau of Reclamation, Technical Service Center, Denver, Colorado, and Albuquerque Area Office, Albuquerque, New Mexico.
- Siegle, R., D. Ahlers, G. Reed, and D. Moore. 2015(a). Bosque del Apache Sediment Plug Baseline Studies; Annual Report 2015. U.S. Department of the Interior, Bureau of Reclamation, Technical Service Center, Denver, Colorado.

- Siegle, R., D. Ahlers, G. Reed, and D. Moore. 2015(b). Elephant Butte Sediment Plug Baseline Studies; Annual Report 2015. U.S. Department of the Interior, Bureau of Reclamation, Technical Service Center, Denver, Colorado.
- Sogge, M.K., Ahlers, D., and Sferra, S.J. 2010. A Natural History Summary and Survey Protocol for the Southwestern Willow Flycatcher: U.S. Geological Survey Techniques and Methods 2A-10, 38 p.
- Sprenger, M.D. 1999. Restoration of Riparian Wildlife Habitat in the Middle Rio Grande Valley Following Historical River Hydrographs. A Thesis In Wildlife Science, Master of Science, Texas Tech University.
- Stoleson, S.H., and D.N. Finch. 1999. Reproductive success of Southwestern Willow Flycatchers in the Cliff Gila Valley, New Mexico. USDA Forest Service Rocky Mountain Research Station report to Phelps Dodge Corporation.
- Stromberg, J.C. 1993. Fremont Cottonwood-Goodding Willow Riparian Forests: A Review of Their Ecology, Threats, and Recovery Potential. Journal of the Arizona-Nevada Academy of Science 27: 97-110.
- USDA, NRCS (U.S. Department of Agriculture, Natural Resources Conservation Service). 1998. 1998 Annual Interagency Riparian Report. Plant Materials Center, Los Lunas, New Mexico.
- USDA, NRCS. 2014. The PLANTS Database (<http://plants.usda.gov>, Accessed 18 March 2014). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.
- USFWS (U.S. Fish and Wildlife Service). 1997. Endangered and Threatened Wildlife and Plants; Final Determination of Critical Habitat for the Southwestern Flycatcher. Federal Register. July 22, 1997. Volume 62, Number 140.
- USFWS. 2001. Programmatic Biological Opinion on the Effects of the U.S. Bureau of Reclamation's, U.S. Army Corps Of Engineers', and Non-Federal Entities' Discretionary Actions Related to Water Management on The Middle Rio Grande, Albuquerque, NM. June 29, 2001.
- USFWS. 2013. Endangered and Threatened Wildlife and Plants; Proposed Threatened Status for the Western Distinct Population Segment of the Yellow-billed Cuckoo (*Coccyzus americanus*); proposed Rule. Federal Register 78(192): 61622.

Appendix A

Waypoint Locations for Avian Point Counts,
Vegetation Transects, Groundwater Monitoring Wells,
and Photo Stations

All coordinates are in NAD83, Zone 13

Avian Point Count Waypoints

Point	Easting (X)	Northing (Y)
LL1-01	340970	3848075
LL1-02	340874	3847961
LL1-03	340818	3847867
LL1-04	340717	3847768
LL1-05	340649	3847675
LL1-06	340612	3847536
LL1-07	340505	3847477
LL1-08	340395	3847340
LL1-09	340410	3847172
LL1-10	340345	3847004
LL1-11	340316	3846827
LL1-12	340267	3846641
LL2-01	341046	3847985
LL2-02	340969	3847883
LL2-03	340900	3847777
LL2-04	340833	3847665
LL2-05	340766	3847559
LL2-06	340696	3847442
LL2-07	340630	3847332
LL2-08	340558	3847202
LL2-09	340502	3847081
LL2-10	340454	3846973
LL2-11	340418	3846865
LL2-12	340380	3846720

Vegetation Transect Waypoints

Transect	x	y
R1A	341053	3847958
R1B	341015	3847992
R2A	340981	3847867
R2B	340943	3847895
R3A	340923	3847761
R3B	340880	3847789
R4A	340860	3847665
R4B	340814	3847687
R5A	340793	3847560
R5B	340749	3847584
R6A	340734	3847459
R6B	340691	3847484
R6-1A	340674	3847363
R6-1B	340630	3847384
R7A	340563	3847162
R7B	340508	3847180
R8A	340516	3847052
R8B	340465	3847073
R9A	340466	3846945
R9B	340417	3846961
R10A	340424	3846834
R10B	340374	3846842
R11A	340392	3846715
R11B	340342	3846723

Groundwater Well Waypoints

Well	x	y
N1	341087	3847987
N2	341037	3848047
N3	340992	3848103
N4	340933	3848162
M1	340613	3847298
M2	340592	3847425
M3	340529	3847439
M4	340469	3847513
S1	340324	3846590
S2	340280	3846598
S3	340245	3846598

Photo Station Waypoints

Photo Station	x	y
P-1	341038	3848023
P-2	340771	3847679
P-3	340582	3847349
P-4	340419	3847015
P-5	340345	3846598
P-6	340898	3848173
P-7	340416	3847477
P-8	340404	3847462
P-9	340384	3847449
P-10	340200	3846582

Appendix B

Bird Species Detected During Point Counts and Associated Habitat Guilds

Species code	Species	Scientific name	Canopy	Cavity	Dense shrub	Edge	Ground shrub	Mid-story	Open-ing	Urban	Water	Migrant
AMAV	American avocet	<i>Recurvirostra americana</i>									X	
AMCR	American crow	<i>Corvus brachyrhynchos</i>				X						
AMPI	American pipit	<i>Anthus rubescens</i>					X					
AM KE	American kestrel	<i>Falco sparverius sparverius</i>		X								
AMRO	American robin	<i>Turdus migratorius</i>						X				
ATFL	Ash-throated flycatcher	<i>Myiarchus cinerascens</i>		X								
BAOW	Barn owl	<i>Tyto alba</i>				X						
BARS	Barn swallow	<i>Hirundo rustica</i>							X			
BANS	Bank swallow	<i>Riparia riparia</i>									X	
BEWR	Bewick's wren	<i>Thryomanes bewickii</i>		X								
BLPH	Black phoebe	<i>Sayornis nigricans</i>									X	
BCCH	Black-capped chickadee	<i>Poecile atricapillus</i>		X								
BCHU	Black-chinned hummingbird	<i>Archilochus alexandri</i>				X						
BCNH	Black-crowned night heron	<i>Nycticorax nycticorax</i>									X	
BHGR	Black-headed grosbeak	<i>Pheucticus melanocephalus</i>						X				
BNST	Black-necked stilt	<i>Himantopus mexicanus</i>									X	
BLGR	Blue grosbeak	<i>Guiraca caerulea</i>					X					
BGGN	Blue-gray gnatcatcher	<i>Poliopitila caerulea</i>						X				
BWTE	Blue-winged teal	<i>Anas discors</i>									X	
BRBL	Brewer's blackbird	<i>Euphagus cyanocephalus</i>										X
BTHU	Broadtailed hummingbird	<i>Selasphorus platycercus</i>										X
BHCO	Brown-headed cowbird	<i>Molothrus ater</i>						X				
BUOR	Bullock's oriole	<i>Icterus bullockii</i>	X									
BUSH	Bushtit	<i>Psaltiriparus minimus</i>						X				
CAGO	Canada goose	<i>Branta canadensis</i>									X	
CAFI	Cassin's finch	<i>Carpodacus cassinii</i>										X
CAVI	Cassin's vireo	<i>Vireo cassinii</i>										X
CAEG	Cattle egret	<i>Bubulcus ibis</i>										X
CLSW	Cliff swallow	<i>Petrochelidon pyrrhonota</i>									X	
COGR	Common grackle	<i>Quiscalus quiscula</i>				X						
COYE	Common yellowthroat	<i>Geothlypis trichas</i>			X							
COHA	Cooper's hawk	<i>Accipiter cooperii</i>	X									

Species code	Species	Scientific name	Canopy	Cavity	Dense shrub	Edge	Ground shrub	Mid-story	Opening	Urban	Water	Migrant
DOWO	Downy woodpecker	<i>Picoides pubescens</i>		X								
DUFL	Dusky flycatcher	<i>Empidonax oberholseri</i>										X
EUST	European starling	<i>Sturnus vulgaris</i>		X								
GADW	Gadwall	<i>Anas strepera</i>										X
GAQU	Gambel's quail	<i>Callipepla gambelii</i>					X					
GRCA	Gray catbird	<i>Dumetella carolinensis</i>						X				
GREG	Great egret	<i>Ardea alba</i>										X
GBHE	Great-blue heron	<i>Ardea herodias</i>									X	
GHOW	Great-horned owl	<i>Bubo virginianus</i>	X									
GTGR	Great-tailed grackle	<i>Quiscalus mexicanus</i>									X	
GRHE	Green heron	<i>Butorides virescens</i>									X	
GTTO	Green-tailed towhee	<i>Pipilo chlorurus</i>					X					
HAWO	Hairy woodpecker	<i>Picoides villosus</i>		X								
HAFL	Hammond's flycatcher	<i>Empidonax hammondi</i>										X
HOFI	House finch	<i>Carpodacus mexicanus</i>						X				
INBU	Indigo bunting	<i>Passerina cyanea</i>				X						
KILL	Killdeer	<i>Charadrius vociferus</i>					X					
LBWO	Ladder-backed woodpecker	<i>Picoides scalaris</i>		X								
LASP	Lark sparrow	<i>Chondestes grammacus</i>					X					
LABU	Lazuli bunting	<i>Passerina amoena</i>										X
LEGO	Lesser goldfinch	<i>Carduelis psaltria</i>						X				
LBHE	Little blue heron	<i>Egretta caerulea</i>										X
LOSH	Loggerhead shrike	<i>Lanius ludovicianus</i>				X						
LUWA	Lucy's warbler	<i>Vermivora luciae</i>										X
MGWA	MacGillivray's warbler	<i>Ardea alba</i>										X
MALL	Mallard	<i>Anas platyrhynchos</i>									X	
MOCH	Mountain chickadee	<i>Poecile gambeli</i>		X								
MODO	Mourning dove	<i>Zenaida macroura</i>					X					
NOFL	Northern flicker	<i>Colaptes auratus</i>		X								
NOMO	Northern mockingbird	<i>Mimus polyglottos</i>				X						
NRWS	Northern rough-winged swallow	<i>Stelgidopteryx serripennis</i>									X	

Species code	Species	Scientific name	Canopy	Cavity	Dense shrub	Edge	Ground shrub	Mid-story	Opening	Urban	Water	Migrant
OCWA	Orange-crowned warbler	<i>Vermivora celata</i>					X					
PHAI	Phainopepla	<i>Phainopepla nitens</i>										X
PLVI	Plumbeous vireo	<i>Vireo plumbeus</i>						X				
RTHA	Red-tailed hawk	<i>Buteo jamaicensis</i>	X									
RWBL	Red-winged blackbird	<i>Agelaius phoeniceus</i>									X	
RNPH	Ring-necked pheasant	<i>Phasianus colchicus</i>					X					
ROPI	Rock pigeon	<i>Columba livia</i>								X		
SAPH	Say's phoebe	<i>Sayornis saya</i>				X						
SNEG	Snowy egret	<i>Egretta thula</i>									X	
SWFL	Southwestern willow flycatcher	<i>Empidonax traillii</i>						X				
SPSA	Spotted sandpiper	<i>Actitis macularia</i>									X	
SPTO	Spotted towhee	<i>Pipilo maculatus</i>						X				
SUTA	Summer tanager	<i>Piranga rubra</i>	X									
SWHA	Swainson's hawk	<i>Buteo swainsoni</i>	X									
TOWA	Townsend's warbler	<i>Dendroica townsendi</i>										X
TRES	Tree swallow	<i>Tachycineta bicolor</i>		X								
TUVU	Turkey vulture	<i>Cathartes aura</i>	X									
UNSW	Unidentified swallow										X	
VGSW	Violet-green swallow	<i>Tachycineta thalassina</i>									X	
WAVI	Warbling vireo	<i>Vireo gilvus</i>						X				
WEKI	Western kingbird	<i>Tyrannus verticalis</i>				X						
WESO	Western screech owl	<i>Otus kennicottii</i>		X								
WETA	Western tanager	<i>Piranga ludoviciana</i>	X									
WEWP	Western wood pewee	<i>Contopus sordidulus</i>	X									
WBNU	White-breasted nuthatch	<i>Sitta carolinensis</i>		X								
WWDO	White-winged dove	<i>Zenaida asiatica</i>						X				
WIWA	Wilson's warbler	<i>Wilsonia pusilla</i>										X
YWAR	Yellow warbler	<i>Dendroica petechia</i>			X							
YBCH	Yellow-breasted chat	<i>Icteria virens</i>						X				
YRWA	Yellow-rumped warbler	<i>Dendroica coronata</i>						X				

Appendix C

Relative Abundance of Individual Bird Species by Area

Table C-1.—Relative abundance of individual bird species in the Cleared/overbank area from 2003 to 2009.

Cleared/overbank area	2003 n=24		2004 n=24		2005 n=24		2006 n=24		2007 n=36		2008 n=36		2009 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Canopy birds														
Turkey vulture	4.2	0.42 (2.04)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Cavity birds														
American kestrel	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Ash-throated flycatcher	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Bewick's wren	0.0	0.00 (0.00)	8.3	0.13 (0.45)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Downy woodpecker	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Northern flicker	0.0	0.00 (0.00)	4.2	0.04 (0.20)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	5.6	0.06 (0.23)	0.0	0.00 (0.00)
White-breasted nuthatch	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.08 (0.28)	0.0	0.00 (0.00)
Dense shrub birds														
Common yellowthroat	0.0	0.00 (0.00)	12.5	0.13 (0.34)	16.7	0.21 (0.51)	16.7	0.17 (0.38)	61.1	0.81 (0.86)	36.1	0.42 (0.60)	47.2	0.50 (0.56)
Edge birds														
American crow	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.21 (1.02)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Black-chinned hummingbird	4.2	0.08 (0.41)	8.3	0.08 (0.28)	12.5	0.13 (0.34)	29.2	0.33 (0.56)	38.9	0.58 (0.84)	33.3	0.47 (0.77)	33.3	0.36 (0.54)
Indigo bunting	8.3	0.08 (0.28)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Loggerhead shrike	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Northern mockingbird	0.0	0.00 (0.00)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	29.2	0.38 (0.71)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Say's phoebe	8.3	0.13 (0.45)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Western kingbird	12.5	0.21 (0.59)	25.0	0.29 (0.55)	16.7	0.21 (0.51)	37.5	0.58 (0.88)	16.7	0.36 (0.90)	2.8	0.03 (0.17)	5.5	0.11 (0.46)
Ground shrub birds														
Blue grosbeak	20.8	0.33 (0.70)	2.1	0.29 (0.62)	4.2	0.04 (0.20)	25.0	0.46 (0.93)	44.4	0.69 (0.89)	13.9	0.14 (0.35)	13.9	0.17 (0.45)
Killdeer	8.3	0.08 (0.28)	37.5	0.67 (1.20)	37.5	0.96 (1.60)	20.8	0.25 (0.53)	22.2	0.42 (0.94)	5.6	0.08 (0.37)	8.3	0.17 (0.56)
Mourning dove	0.0	0.00 (0.00)	16.7	0.17 (0.38)	12.5	0.25 (0.74)	45.8	3.92 (7.63)	25.0	0.69 (2.08)	19.4	0.28 (0.66)	25.0	0.42 (0.87)
Ring-necked pheasant	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.08 (0.28)	14.9	0.14 (0.35)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Midstory birds														
American robin	0.0	0.00 (0.00)	4.2	0.04 (0.20)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Black-headed grosbeak	4.2	0.04 (0.20)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	19.4	0.28 (0.61)	22.2	0.22 (0.42)
Brown-headed cowbird	8.3	0.08 (0.28)	29.2	0.54 (0.98)	0.0	0.00 (0.00)	12.5	0.25 (0.68)	25.0	0.50 (1.00)	50.0	1.17 (1.75)	8.3	0.17 (0.61)
Bushtit	0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.00	0.00 (0.00)	2.8	0.11 (0.67)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Gray catbird	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.04 (0.20)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	2.8	0.03 (0.17)
House finch	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.13 (0.61)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Lesser goldfinch	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Spotted towhee	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.00	0.00 (0.00)	16.7	0.19 (0.47)	25.0	0.28 (0.51)	33.3	0.39 (0.60)
Southwestern willow flycatcher	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.00	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
White-winged dove	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.00	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	2.8	0.03 (0.17)

<i>Cleared/overbank area</i>	2003 n=24		2004 n=24		2005 n=24		2006 n=24		2007 n=36		2008 n=36		2009 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Yellow-breasted chat	0.0	0.00 (0.00)	4.2	0.04 (0.20)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	13.9	0.17 (0.45)	2.8	0.03 (0.17)
Open birds														
Barn swallow	4.2	0.08 (0.41)	16.7	0.17 (0.38)	8.3	0.08 (0.28)	2.1	0.58 (1.32)	2.8	0.11 (0.67)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Water birds														
American avocet	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Black-crowned night heron	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	8.3	0.11 (0.40)	0.0	0.00 (0.00)
Black-necked stilt	0.0	0.00 (0.00)	4.2	0.17 (0.82)	25.0	0.42 (0.83)	8.3	0.13 (0.45)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Blue-winged teal	0.0	0.00 (0.00)	0.0	0.00 (0.00)	12.5	0.21 (0.66)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Cliff swallow	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.17 (0.61)	0.0	0.00 (0.00)
Great-blue heron	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.5	0.06 (0.23)
Great-tailed grackle	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Mallard	0.0	0.00 (0.00)	0.0	0.00 (0.00)	33.3	1.46 (3.16)	4.2	0.04 (0.20)	5.6	0.11 (0.52)	8.3	0.11 (0.40)	5.5	0.06 (0.23)
Northern rough-winged swallow	12.5	0.13 (0.34)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.17 (0.61)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Red-winged blackbird	4.2	0.67 (1.13)	50.0	1.21 (1.50)	95.8	4.63 (1.79)	33.3	0.46 (0.78)	47.2	1.11 (1.69)	55.6	1.28 (1.60)	41.7	0.58 (0.81)
Snowy egret	12.5	0.13 (0.34)	20.8	0.29 (0.62)	12.5	0.21 (0.59)	0.0	0.00 (0.00)	8.3	0.11 (0.40)	5.6	0.06 (0.23)	2.8	0.03 (0.17)
Spotted sandpiper	12.5	0.13 (0.34)	12.5	0.17 (0.48)	37.5	0.46 (0.66)	8.3	0.13 (0.45)	8.3	0.08 (0.28)	5.6	0.08 (0.37)	0.0	0.00 (0.00)
Unidentified swallow	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	25.0	0.33 (0.64)	2.8	0.08 (0.50)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Violet-green swallow	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	25.0	0.38 (0.71)	2.8	0.03 (0.17)	8.3	0.17 (0.61)	0.0	0.00 (0.00)
Migrants														
Brewer's blackbird	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.00	0.00 (0.00)	0.00	0.00 (0.00)	2.8	1.25 (7.50)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Cassin's finch	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Cattle egret	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.25 (1.22)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Gadwall	0.0	0.00 (0.00)	0.0	0.00 (0.00)	4.2	0.13 (0.61)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Lazuli bunting	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.11 (0.40)	0.0	0.00 (0.00)
Little blue heron	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Lucy's warbler	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)

Table C-1.(cont'd)—Relative abundance of individual bird species in the Cleared/overbank area from 2010 to 2016.

<i>Cleared/overbank area</i>	2010 n=36		2011 n=36		2012 n=36		2013 n=36		2014 n=36		2015 n=36		2016 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Canopy birds														
Bullock's oriole	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	11.1	0.11 (0.32)	2.8	0.03 (0.17)
Cooper's hawk	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	5.6	0.06 (0.23)
Summer tanager	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	11.1	0.11 (0.32)	13.9	0.14 (0.35)
Swainson's hawk	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	2.8	0.03 (0.17)

Cleared/overbank area	2010 n=36		2011 n=36		2012 n=36		2013 n=36		2014 n=36		2015 n=36		2016 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Western tanager	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Western wood pewee	0.0	0.00 (0.00)	0.0	0.03 (0.17)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Cavity birds														
American kestrel	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)
Ash-throated flycatcher	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)	2.8	0.03 (0.17)	13.9	0.14 (0.35)	5.6	0.06 (0.23)	13.9	0.06 (0.23)
Bewick's wren	0.0	0.00 (0.00)	16.7	0.22 (0.54)	2.8	0.03 (0.17)	8.3	0.08 (0.28)	2.8	0.06 (0.33)	11.1	0.11 (0.32)	16.7	0.19 (0.47)
Black-capped chickadee	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Downy woodpecker	0.0	0.00 (0.00)	0.0	0.00 (0.00)	13.9	0.22 (0.59)	2.8	0.06 (0.33)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Ladder-backed woodpecker	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.06 (0.23)
Mountain chickadee	8.3	0.11 (0.40)	8.3	0.08 (0.28)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.06 (0.33)
Northern flicker	8.3	0.08 (0.28)	2.8	0.03 (0.17)	5.6	0.06 (0.23)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	8.3	0.11 (0.40)
Western screech-owl	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Dense shrub birds														
Common yellowthroat	25.0	0.25 (0.44)	47.2	0.56 (0.65)	41.7	0.58 (0.77)	11.1	0.11 (0.32)	22.2	0.28 (0.57)	47.2	0.56 (0.69)	19.4	0.28 (0.61)
Yellow warbler	2.8	0.03 (0.17)	5.5	0.06 (0.23)	5.6	0.06 (0.23)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	8.3	0.11 (0.40)	0.0	0.00 (0.00)
Edge birds														
Black-chinned hummingbird	44.4	0.53 (0.65)	41.7	0.56 (0.73)	55.6	0.83 (0.85)	66.7	0.92 (0.77)	72.2	1.28 (1.11)	44.4	0.67 (0.86)	33.3	0.44 (0.73)
Indigo bunting	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.08 (0.28)	8.3	0.08 (0.28)
Northern mockingbird	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Say's phoebe	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.11 (0.46)	2.8	0.03 (0.17)	5.6	0.06 (0.23)	2.8	0.03 (0.17)
Western kingbird	2.8	0.03 (0.17)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Ground shrub birds														
American pipit	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Blue grosbeak	11.1	0.11 (0.32)	25.0	0.33 (0.63)	13.9	0.22 (0.59)	22.2	0.33 (0.72)	19.4	0.25 (0.55)	27.8	0.39 (0.69)	27.8	0.36 (0.64)
Killdeer	5.6	0.11 (0.52)	8.3	0.11 (0.40)	8.3	0.08 (0.28)	19.4	0.31 (0.71)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Mourning dove	25.0	0.33 (0.63)	36.1	0.53 (0.84)	55.6	0.86 (0.87)	55.6	0.78 (0.80)	55.6	0.83 (0.88)	25.0	0.31 (0.58)	50.0	0.75 (0.87)
Orange-crowned warbler	0.00	0.00 (0.00)	0.00	0.00 (0.00)	13.9	0.17 (0.45)	13.9	0.25 (0.65)	11.1	0.14 (0.42)	8.3	0.08 (0.28)	5.6	0.08 (0.37)
Ring-necked pheasant	2.8	0.03 (0.17)	8.3	0.08 (0.28)	22.2	0.22 (0.42)	19.4	0.19 (0.40)	16.7	0.17 (0.38)	8.3	0.08 (0.28)	8.3	0.08 (0.28)
Midstory birds														
American robin	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	13.9	0.19 (0.52)	11.1	0.11 (0.32)
Black-headed grosbeak	33.3	0.50 (0.81)	38.9	0.50 (0.70)	66.7	0.92 (0.77)	61.1	0.75 (0.69)	50.0	0.64 (0.72)	44.4	0.61 (0.77)	41.7	0.61 (0.87)
Blue-gray gnatcatcher	0.0	0.00 (0.00)	5.5	0.08 (0.37)	5.6	0.08 (0.37)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Brown-headed cowbird	36.1	0.61 (0.96)	41.7	0.78 (1.07)	66.7	1.28 (1.21)	58.3	1.03 (1.16)	41.7	0.67 (0.93)	16.7	0.36 (1.10)	11.1	0.14 (0.42)
Bushtit	2.8	0.17 (1.00)	5.5	0.14 (0.59)	8.3	0.25 (0.84)	11.1	0.25 (0.77)	5.6	0.17 (0.74)	2.8	0.03 (0.17)	5.6	0.08 (0.37)
Gray catbird	5.6	0.06 (0.23)	2.8	0.03 (0.17)	5.6	0.06 (0.23)	5.6	0.06 (0.23)	11.1	0.14 (0.42)	25.0	0.28 (0.51)	30.6	0.47 (0.81)

Cleared/overbank area	2010 n=36		2011 n=36		2012 n=36		2013 n=36		2014 n=36		2015 n=36		2016 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
House finch	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)	2.8	0.03 (0.17)	8.3	0.17 (0.56)	13.9	0.19 (0.52)	13.9	0.14 (0.35)
Lesser goldfinch	0.0	0.00 (0.00)	5.5	0.14 (0.59)	25.0	0.47 (0.88)	8.3	0.17 (0.56)	8.3	0.14 (0.54)	2.8	0.03 (0.17)	8.3	0.11 (0.40)
Plumbeous vireo	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Spotted towhee	55.6	0.64 (0.64)	41.7	0.50 (0.65)	66.7	1.06 (0.89)	94.4	1.31 (0.58)	69.4	1.03 (0.84)	63.9	0.81 (0.71)	88.9	1.76 (0.96)
White-winged dove	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)	5.6	0.08 (0.37)
Yellow-breasted chat	5.6	0.06 (0.23)	80.5	1.19 (0.79)	75.0	1.17 (0.85)	91.7	1.61 (0.80)	75.0	1.31 (0.95)	47.2	0.53 (0.61)	41.7	0.56 (0.73)
Yellow-rumped warbler	0.0	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Open birds														
Barn swallow	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	25.0	0.81 (1.74)	2.8	0.06 (0.33)
Water birds														
Bank swallow	0.0	0.00 (0.00)	2.8	0.08 (0.50)	0.0	0.00 (0.00)	2.8	0.11 (0.67)	0.0	0.00 (0.00)	13.9	0.33 (0.86)	0.0	0.00 (0.00)
Black-crowned night heron	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)
Black phoebe	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Canada goose	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.31 (1.09)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Great-blue heron	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Green heron	0.0	0.00 (0.00)	5.5	0.06 (0.23)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	2.8	0.03 (0.17)
Mallard	2.8	0.22 (1.33)	11.1	0.31 (1.09)	5.6	0.08 (0.37)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.11 (0.40)	5.6	0.06 (0.23)
Red-winged blackbird	8.3	0.17 (0.70)	11.1	0.22 (0.64)	5.6	0.14 (0.59)	8.3	0.14 (0.49)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Snowy egret	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Migrants														
Broadtailed hummingbird	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	41.7	0.78 (1.02)
Cassin's vireo	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Dusky flycatcher	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Hammond's flycatcher	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Great egret	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
MacGillivray's warbler	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Phainopepla	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)	0.0	0.00 (0.00)
Townsend's warbler	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Wilson's warbler	0.0	0.00 (0.00)	8.3	0.08 (0.28)	11.1	0.11 (0.32)	8.3	0.11 (0.40)	8.3	0.11 (0.40)	2.8	0.03 (0.17)	5.6	0.06 (0.24)

Table C-2.—Relative abundance of individual bird species in the Burned area in 2003 and 2004 and 2007 to 2010.

Burned area	2003 n=42		2004 n=47		2007 n=36		2008 n=36		2009 n=36		2010 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Canopy birds												
Cooper's hawk	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.08 (0.28)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	0.0	0.00 (0.00)
Great-horned owl	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Red-tailed hawk	4.8	0.05 (0.22)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Summer tanager	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.08 (0.28)	0.0	0.00 (0.00)
Turkey vulture	19.0	0.67 (1.72)	8.5	0.36 (1.28)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Western tanager	2.4	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Western wood pewee	0.0	0.00 (0.00)	2.1	0.02 (0.15)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	0.0	0.00 (0.00)
Cavity birds												
American kestrel	7.1	0.10 (0.37)	2.1	0.02 (0.15)	13.9	0.17 (0.45)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)
Ash-throated flycatcher	19.0	0.19 (0.40)	6.4	0.06 (0.25)	11.1	0.14 (0.42)	2.8	0.03 (0.17)	8.3	0.08 (0.28)	11.1	0.11 (0.32)
Bewick's wren	4.8	0.05 (0.22)	0.0	0.00 (0.00)	25.0	0.39 (0.80)	5.6	0.06 (0.23)	8.3	0.08 (0.28)	13.9	0.17 (0.45)
Black-capped chickadee	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Downy woodpecker	0.0	0.00 (0.00)	2.1	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
European starling	2.4	0.02 (0.15)	2.1	0.02 (0.15)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Hairy woodpecker	0.0	0.00 (0.00)	4.3	0.04 (0.20)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Ladder-backed woodpecker	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.08 (0.28)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Northern flicker	19.0	0.21 (0.47)	10.6	0.11 (0.31)	22.2	0.25 (0.50)	5.6	0.06 (0.23)	16.7	0.19 (0.37)	8.3	0.11 (0.40)
White-breasted nuthatch	7.1	0.07 (0.26)	17.0	0.17 (0.38)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Dense shrub birds												
Common yellowthroat	19.0	0.19 (0.40)	10.6	0.11 (0.31)	16.7	0.17 (0.38)	13.9	0.14 (0.35)	2.8	0.03 (0.17)	2.8	0.03 (0.17)
Yellow warbler	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Edge birds												
Black-chinned hummingbird	45.2	0.57 (0.74)	46.8	0.51 (0.59)	75.0	1.08 (0.81)	44.4	0.44 (0.50)	77.8	1.28 (0.88)	77.8	1.31 (1.09)
Common grackle	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Northern mockingbird	2.4	0.05 (0.31)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Say's phoebe	2.4	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Western kingbird	11.9	0.19 (0.59)	17.0	0.19 (0.45)	30.6	0.56 (0.91)	5.6	0.06 (0.23)	5.6	0.11 (0.46)	5.6	0.08 (0.37)
Ground shrub birds												
Blue grosbeak	33.3	0.40 (0.63)	21.3	0.26 (0.53)	8.3	0.11 (0.40)	11.1	0.11 (0.32)	2.8	0.06 (0.33)	2.8	0.03 (0.17)
Gambel's quail	0.0	0.00 (0.00)	2.1	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)

Burned area	2003 n=42		2004 n=47		2007 n=36		2008 n=36		2009 n=36		2010 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Killdeer	2.4	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Mourning dove	4.8	0.67 (0.90)	61.7	0.96 (0.88)	58.3	1.36 (1.64)	44.4	0.61 (0.80)	38.9	0.64 (0.99)	38.9	0.58 (0.81)
Ring-necked pheasant	4.8	0.05 (0.22)	4.2	0.04 (0.20)	16.7	0.28 (0.78)	13.9	0.14 (0.35)	16.7	0.17 (0.38)	19.4	0.22 (0.48)
Midstory birds												
American robin	4.8	0.05 (0.22)	14.9	0.21 (0.59)	8.3	0.08 (0.28)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	5.6	0.08 (0.37)
Black-headed grosbeak	69.0	1.00 (0.88)	61.7	0.74 (0.67)	44.4	0.56 (0.81)	58.3	0.83 (0.85)	47.2	0.69 (0.89)	41.7	0.53 (0.70)
Blue-gray gnatcatcher	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)
Brown-headed cowbird	66.7	1.36 (1.43)	36.2	0.66 (1.13)	58.3	0.86 (0.96)	55.6	0.92 (1.34)	36.1	0.64 (0.99)	27.8	0.53 (1.03)
Bushtit	0.0	0.00 (0.00)	2.1	0.11 (0.73)	5.6	0.17 (0.85)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Gray catbird	26.2	0.26 (0.45)	48.9	0.53 (0.58)	36.1	0.50 (0.74)	22.2	0.28 (0.57)	50.0	0.77 (0.76)	44.4	0.56 (0.69)
House finch	2.4	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Lesser goldfinch	2.4	0.05 (0.31)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Spotted towhee	50.0	0.69 (0.84)	80.8	0.91 (0.54)	61.1	0.94 (0.89)	41.7	0.44 (0.56)	41.7	0.56 (0.73)	44.4	0.58 (0.77)
White-winged dove	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Yellow-breasted chat	76.2	1.26 (0.91)	70.2	1.13 (1.03)	38.9	0.47 (0.70)	41.7	0.44 (0.56)	41.7	0.47 (0.61)	30.6	0.33 (0.53)
Open birds												
Barn swallow	2.4	0.02 (0.15)	2.1	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Water birds												
Black phoebe	0.0	0.00 (0.00)	2.1	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Black-necked stilt	2.4	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Red-winged blackbird	9.5	0.12 (0.40)	6.4	0.06 (0.25)	16.7	0.42 (1.16)	11.1	0.69 (2.36)	11.1	0.22 (0.76)	5.5	0.14 (0.68)
Snowy egret	0.0	0.00 (0.00)	2.1	0.02 (0.15)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Spotted sandpiper	4.8	0.05 (0.22)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Migrants												
Lazuli bunting	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)	0.0	0.00 (0.00)	0.0	0.00 (0.00)

Table C-2(cont'd) .—Relative abundance of individual bird species in the Burned area from 2011 to 2015.

Burned area	2011 n=36		2012 n=36		2013 n=36		2014 n=36		2015 n=36		2016 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Canopy birds												
Bullock's oriole	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.7	0.03 (0.17)	11.1	0.03 (0.17)	11.1	0.14 (0.42)	0.0	0.00 (0.00)
Cooper's hawk	0.0	0.00 (0.00)	5.6	0.06 (0.23)	8.3	0.11 (0.40)	2.8	0.03 (0.17)	5.6	0.06 (0.23)	5.6	0.11 (0.52)
Summer tanager	5.5	0.06 (0.23)	8.3	0.08 (0.28)	8.3	0.11 (0.40)	5.6	0.06 (0.23)	2.8	0.03 (0.17)	11.1	0.14 (0.42)

Burned area	2011 n=36		2012 n=36		2013 n=36		2014 n=36		2015 n=36		2016 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Swainson's hawk	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Western wood pewee	5.5	0.06 (0.23)	2.8	0.03 (0.17)	0.0	0.03 (0.17)	2.8	0.03 (0.17)	5.6	0.06 (0.23)	2.8	0.03 (0.17)
Cavity birds												
American kestrel	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Ash-throated flycatcher	8.3	0.08 (0.28)	22.2	0.25 (0.50)	16.7	0.19 (0.47)	8.3	0.08 (0.28)	5.6	0.06 (0.23)	11.1	0.11 (0.32)
Bewick's wren	13.9	0.25 (0.73)	19.4	0.33 (0.76)	11.1	0.11 (0.32)	2.8	0.03 (0.17)	22.2	0.28 (0.57)	25.0	0.25 (0.44)
Black-capped chickadee	0.0	0.00 (0.00)	5.6	0.08 (0.37)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Downy woodpecker	0.0	0.00 (0.00)	5.6	0.06 (0.23)	8.3	0.14 (0.49)	8.3	0.11 (0.40)	2.8	0.03 (0.17)	0.0	0.03 (0.17)
Mountain chickadee	11.1	0.14 (0.42)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	8.3	0.17 (0.70)
Northern flicker	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	2.8	0.03 (0.17)
Western screech-owl	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
White-breasted nuthatch	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)	2.8	0.06 (0.33)
Dense shrub birds												
Common yellowthroat	13.9	0.17 (0.45)	8.3	0.14 (0.49)	0.0	0.00 (0.00)	19.4	0.22 (0.48)	30.6	0.31 (0.47)	16.7	0.17 (0.38)
Yellow warbler	5.5	0.06 (0.23)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Edge birds												
American crow	0.0	0.00 (0.00)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Barn owl	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Black-chinned hummingbird	83.3	1.00 (0.59)	83.3	1.14 (0.76)	91.7	1.47 (0.74)	75.0	1.31 (1.01)	75.0	0.97 (0.70)	39.8	0.64 (0.96)
Indigo bunting	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	8.3	0.08 (0.28)	2.8	0.03 (0.17)
Loggerhead shrike	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Northern mockingbird	5.5	0.06 (0.23)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Say's phoebe	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Western kingbird	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Ground shrub birds												
Blue grosbeak	19.4	0.25 (0.55)	2.8	0.03 (0.17)	22.2	0.25 (0.50)	16.7	0.25 (0.60)	22.2	0.25 (0.50)	16.7	0.28 (0.78)
Gambel's quail	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	5.6	0.08 (0.37)
Mourning dove	55.5	1.03 (1.08)	72.2	1.00 (0.79)	58.3	0.86 (0.87)	69.4	1.00 (0.86)	36.1	0.44 (0.69)	47.2	0.67 (0.83)
Orange-crowned warbler	2.8	0.06 (0.33)	8.3	0.08 (0.28)	16.7	0.19 (0.47)	2.8	0.06 (0.33)	5.6	0.06 (0.23)	0.0	0.00 (0.00)
Ring-necked pheasant	5.5	0.06 (0.23)	16.7	0.17 (0.38)	2.8	0.03 (0.17)	13.9	0.14 (0.35)	5.6	0.06 (0.23)	19.4	0.19 (0.40)
Midstory birds												
American robin	2.8	0.06 (0.33)	8.3	0.08 (0.28)	13.9	0.14 (0.35)	0.0	0.00 (0.00)	8.3	0.08 (0.28)	19.4	0.25 (0.55)
Black-headed grosbeak	36.1	0.47 (0.70)	55.6	0.75 (0.77)	38.9	0.58 (0.81)	47.2	0.69 (0.82)	30.6	0.42 (0.69)	69.4	1.03 (0.88)

Burned area	2011 n=36		2012 n=36		2013 n=36		2014 n=36		2015 n=36		2016 n=36	
Species	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)	% Plots	Mean (SD)
Blue-gray gnatcatcher	5.5	0.06 (0.23)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Brown-headed cowbird	44.4	0.69 (0.92)	25.0	0.42 (0.77)	27.8	0.64 (1.17)	25.0	0.33 (0.63)	8.3	0.11 (0.40)	8.3	0.08 (0.28)
Bushtit	11.1	0.22 (0.68)	5.6	0.08 (0.37)	11.1	0.31 (0.92)	11.1	0.25 (0.81)	11.1	0.25 (0.81)	5.6	0.28 (1.37)
Gray catbird	41.7	0.53 (0.70)	47.2	0.67 (0.79)	44.4	0.61 (0.77)	27.8	0.42 (0.73)	52.8	0.61 (0.69)	25.0	0.33 (0.63)
House finch	5.5	0.17 (0.70)	2.8	0.06 (0.33)	0.0	0.00 (0.00)	2.8	0.06 (0.33)	11.1	0.11 (0.32)	5.6	0.08 (0.37)
Lesser goldfinch	13.9	0.25 (0.69)	5.6	0.08 (0.37)	5.6	0.06 (0.23)	5.6	0.06 (0.23)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Plumbeous vireo	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Spotted towhee	44.4	0.64 (0.80)	55.6	0.78 (0.80)	69.4	0.94 (0.75)	75.0	1.06 (0.79)	47.2	0.61 (0.73)	66.7	1.14 (1.10)
Warbling vireo	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)
White-winged dove	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	16.7	0.17 (0.38)	5.6	0.06 (0.23)
Yellow-breasted chat	72.2	1.06 (0.79)	69.4	1.03 (0.81)	80.6	1.36 (0.87)	88.9	1.61 (0.80)	44.4	0.56 (0.69)	44.4	0.64 (0.83)
Yellow-rumped warbler	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.11 (0.46)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Urban birds												
Rock pigeon	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Water birds												
Black phoebe	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Canada goose	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)
Great-blue heron	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	2.8	0.03 (0.17)
Mallard	5.5	0.22 (1.05)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.06 (0.33)
Red-winged blackbird	11.1	0.28 (0.81)	13.9	0.33 (0.93)	13.9	0.28 (0.74)	8.3	0.14 (0.49)	0.0	0.00 (0.00)	8.3	0.11 (0.40)
Migrants												
Broadtailed hummingbird	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	25.0	0.25 (0.44)	58.3	0.78 (0.80)
Cassin's vireo	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Cattle egret	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Dusky flycatcher	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Lazuli bunting	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.08 (0.37)
MacGillivray's warbler	2.8	0.03 (0.17)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Phainopepla	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Townsend's warbler	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	5.6	0.06 (0.23)	0.0	0.00 (0.00)	0.0	0.00 (0.00)
Wilson's warbler	2.8	0.03 (0.17)	13.9	0.14 (0.34)	0.0	0.00 (0.00)	2.8	0.03 (0.17)	2.8	0.03 (0.17)	0.0	0.00 (0.00)
Black-necked stilt	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)	0.0	0.00 (0.00)

Appendix D

Avian Abundance by Species Guilds

Table D-1.—Total, mean, and standard deviation by species guilds for the Cleared/Overbank Area from 2003 to 2009.

Los Lunas Cleared/overbank area	2003 8 points		2004 8 points		2005 8 points		2006 8 points		2007 12 points		2008 12 points		2009 12 points	
	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)
# Species	18	1.79 (1.25)	20	2.92 (1.61)	21	3.58 (1.35)	20	3.67 (2.04)	24	3.78 (1.66)	22	3.42 (1.71)	18	2.67 (1.45)
# Birds	22	2.75 (3.08)	37	4.58 (2.92)	77	9.67 (4.47)	70	8.79 (9.14)	79	7.83 (11.21)	66	5.50 (3.26)	40	3.36 (2.09)
# Canopy spp.	1	0.04 (0.20)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)
# Canopy birds	3	0.42 (2.04)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)
# Cavity spp.	1	0.04 (0.20)	2	0.13 (0.45)	2	0.08 (0.28)	1	0.04 (0.20)	1	0.06 (0.23)	2	0.14 (0.49)	0	0.00 (0.00)
# Cavity birds	1	0.04 (0.20)	2	0.17 (0.56)	2	0.08 (0.28)	1	0.04 (0.20)	1	0.06 (0.23)	2	0.14 (0.49)	0	0.00 (0.00)
# Dense shrub spp.	0	0.00 (0.00)	1	0.13 (0.34)	1	0.17 (0.38)	1	0.17 (0.38)	1	0.61 (0.49)	1	0.36 (0.49)	1	0.47 (0.51)
# Dense shrub birds	0	0.00 (0.00)	1	0.13 (0.34)	2	0.21 (0.51)	1	0.17 (0.38)	10	0.81 (0.86)	5	0.42 (0.60)	6	0.50 (0.56)
# Edge spp.	5	0.38 (0.65)	5	0.46 (0.59)	2	0.29 (0.46)	4	1.00 (1.06)	3	0.58 (0.65)	2	0.36 (0.49)	2	0.39 (0.55)
# Edge birds	5	0.54 (1.02)	5	0.50 (0.66)	3	0.33 (0.56)	12	1.50 (1.84)	11	2.19 (8.09)	6	0.50 (0.77)	6	0.47 (0.74)
# Ground shrub spp.	2	0.29 (0.46)	3	0.75 (0.79)	3	0.54 (0.59)	4	1.00 (0.83)	4	1.06 (0.89)	4	0.42 (0.60)	3	0.47 (0.70)
# Ground shrub birds	3	0.42 (0.72)	9	1.13 (1.54)	10	1.25 (1.62)	38	4.71 (7.80)	23	1.94 (2.40)	6	0.53 (0.84)	9	0.75 (1.23)
# Mid-story spp.	3	0.17 (0.38)	4	0.42 (0.78)	3	0.13 (0.45)	2	0.17 (0.48)	7	0.61 (0.73)	5	1.11 (0.95)	7	0.75 (0.73)
# Mid-story birds	3	0.17 (0.38)	5	0.67 (1.20)	3	0.21 (0.83)	2	0.29 (0.81)	12	1.00 (1.37)	23	1.92 (1.92)	11	0.89 (0.95)
# Opening spp.	1	0.04 (0.20)	1	0.17 (0.38)	1	0.08 (0.28)	1	0.21 (0.41)	1	0.03 (0.17)	0	0.00 (0.00)	1	0.03 (0.17)
# Opening birds	2	0.08 (0.41)	1	0.17 (0.38)	1	0.08 (0.28)	5	0.58 (1.32)	2	0.11 (0.67)	0	0.00 (0.00)	1	0.03 (0.17)
# Water spp.	5	0.83 (0.83)	4	0.88 (0.90)	9	2.29 (1.08)	7	1.08 (0.83)	8	0.86 (1.05)	8	1.03 (1.06)	4	0.56 (0.73)
# Water birds	9	1.08 (1.21)	15	1.83 (2.48)	60	7.50 (3.88)	12	1.50 (1.25)	20	1.69 (2.25)	24	2.00 (2.07)	8	0.75 (1.05)

Table D-1 (cont'd).—Total, mean, and standard deviation by species guilds for the Cleared/Overbank Area from 2009 to 2016.

Los Lunas Cleared/overbank area	2010 12 points		2011 12 points		2012 12 points		2013 12 points		2014 12 points		2015 12 points		2016 12 points	
	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)
# Species	18	2.86 (1.53)	34	4.86 (1.05)	26	5.89 (1.04)	27	5.92 (1.00)	31	5.44 (1.42)	36	5.33 (0.99)	36	5.17 (1.67)
# Birds	48	4.03 (3.08)	83	6.94 (2.33)	109	9.08 (2.20)	106	8.81 (1.89)	100	8.33 (2.93)	89	7.39 (2.07)	87	7.25 (2.51)
# Canopy spp.	0	0.00 (0.00)	2	0.06 (0.23)	1	0.03 (0.17)	3	0.08 (0.28)	4	0.19 (0.47)	3	0.25 (0.44)	5	0.28 (0.51)
# Canopy birds	0	0.00 (0.00)	2	0.06 (0.23)	1	0.03 (0.17)	3	0.08 (0.28)	4	0.19 (0.47)	3	0.25 (0.44)	5	0.28 (0.51)
# Cavity spp.	2	0.17 (0.45)	4	0.31 (0.52)	4	0.28 (0.45)	4	0.19 (0.47)	4	0.25 (0.44)	5	0.25 (0.44)	5	0.47 (0.70)
# Cavity birds	7	0.19 (0.52)	4	0.36 (0.64)	5	0.39 (0.69)	4	0.22 (0.54)	4	0.31 (0.58)	5	0.28 (0.51)	7	0.56 (0.88)
# Dense shrub spp.	2	0.28 (0.45)	2	0.53 (0.51)	2	0.47 (0.56)	2	0.14 (0.35)	2	0.25 (0.50)	2	0.56 (0.61)	1	0.19 (0.40)
# Dense shrub birds	3	0.28 (0.45)	7	0.61 (0.64)	10	0.81 (0.82)	2	0.14 (0.35)	4	0.31 (0.67)	8	0.67 (0.83)	3	0.28 (0.62)

# Edge spp.	2	0.47 (0.56)	4	0.50 (0.56)	2	0.58 (0.50)	2	0.72 (0.51)	3	0.78 (0.48)	4	0.61 (0.55)	4	0.61 (0.55)
# Edge birds	7	0.56 (0.73)	7	0.64 (0.76)	10	0.86 (0.83)	12	1.03 (0.84)	16	1.33 (1.10)	10	0.83 (0.88)	7	0.58 (0.81)
# Ground shrub spp.	4	0.44 (0.69)	4	0.78 (0.64)	5	1.14 (0.76)	5	1.31 (0.79)	5	1.08 (0.69)	5	0.72(0.66)	4	0.92 (0.73)
# Ground shrub birds	7	0.58 (1.00)	13	1.06 (1.09)	19	1.56 (1.08)	22	1.86 (1.22)	17	1.44 (1.03)	11	0.89 (0.89)	15	1.28 (1.19)
# Mid-story spp.	6	1.39 (0.99)	12	2.33 (0.93)	9	3.25 (0.87)	9	3.36 (0.90)	10	2.75 (1.05)	10	2.36 (0.90)	12	2.67 (1.41)
# Mid-story birds	24	2.03 (1.93)	42	3.50 (1.76)	64	5.36 (1.97)	63	5.22 (1.99)	52	4.36 (2.22)	37	3.11 (1.39)	49	4.10 (2.18)
# Opening spp.	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	1	0.39 (0.49)	1	0.03 (0.17)
# Opening birds	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	10	1.14 (1.79)	1	0.06 (0.33)
# Water spp.	2	0.11 (0.32)	6	0.36 (0.64)	3	0.14 (0.35)	2	0.11 (0.32)	3	0.14 (0.42)	6	0.33 (0.59)	4	0.14 (0.35)
# Water birds	5	0.39 (1.48)	9	0.72 (1.58)	3	0.25 (0.69)	3	0.25 (0.81)	5	0.39 (1.18)	7	0.56 (1.03)	4	0.14 (0.35)

Table D-2.—Total, mean, and standard deviation by species guilds for the Burned Area from 2003 to 2004 and 2007 to 2010.

Los Lunas Burned area	2003 17 points		2004 17 points		2007 12 points		2008 12 points		2009 12 points		2010 12 points	
	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)
# Species	30	5.71 (1.66)	27	5.47 (1.40)	24	5.81 (2.23)	17	3.83 (1.54)	24	4.42 (1.44)	18	3.89 (1.53)
# Birds	146	8.45 (3.23)	118	7.34 (2.55)	107	8.89 (3.77)	65	5.42 (3.55)	75	6.28 (2.35)	66	5.50 (2.81)
# Canopy spp.	3	0.26 (0.50)	2	0.11 (0.31)	2	0.14 (0.35)	0	0.00 (0.00)	4	0.22 (0.42)	0	0.00 (0.00)
# Canopy birds	11	0.74 (1.80)	6	0.38 (1.28)	2	0.14 (0.35)	0	0.00 (0.00)	3	0.22 (0.42)	0	0.00 (0.00)
# Cavity spp.	6	0.60 (0.70)	7	0.45 (0.69)	6	0.83 (0.97)	3	0.14 (0.35)	4	0.36 (0.59)	4	0.39 (0.55)
# Cavity birds	12	0.62 (0.76)	7	0.45 (0.69)	13	1.08 (1.38)	3	0.14 (0.35)	5	0.39 (0.64)	6	0.47 (0.70)
# Dense shrub spp.	1	0.19 (1.40)	1	0.11 (0.31)	1	0.17 (0.38)	1	0.14 (0.35)	2	0.06 (0.23)	1	0.03 (0.17)
# Dense shrub birds	3	0.19 (1.40)	2	0.11 (0.31)	2	0.17 (0.38)	2	0.14 (0.35)	2	0.06 (0.23)	1	0.03 (0.17)
# Edge spp.	4	0.62 (0.58)	2	0.64 (0.61)	3	1.08 (0.65)	3	0.53 (0.70)	3	0.86 (0.42)	2	0.83 (0.51)
# Edge birds	15	0.83 (0.93)	12	0.70 (0.69)	20	1.69 (1.21)	6	0.53 (0.70)	17	1.42 (0.87)	17	1.39 (1.13)
# Ground shrub spp.	4	0.88 (0.80)	4	0.89 (0.70)	3	0.83 (0.61)	3	0.69 (0.71)	3	0.58 (0.60)	3	0.61 (0.65)
# Ground shrub birds	18	1.14 (1.26)	20	1.28 (1.04)	21	1.75 (1.73)	10	0.86 (1.05)	10	0.86 (1.13)	10	0.83 (0.94)
# Mid-story spp.	8	2.98 (1.18)	7	3.15 (0.98)	8	2.58 (1.18)	6	2.22 (1.10)	7	2.22 (1.35)	7	1.97 (1.08)
# Mid-story birds	83	4.69 (2.28)	69	4.30 (1.94)	44	3.64 (1.96)	37	3.06 (1.82)	37	3.11 (2.14)	32	2.64 (1.89)
# Opening spp.	1	0.02 (0.15)	1	0.02 (0.15)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)
# Opening birds	1	0.02 (0.15)	1	0.02 (0.15)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)
# Water spp.	3	0.17 (0.38)	3	0.11 (0.31)	1	0.17 (0.38)	1	0.11 (0.32)	1	0.11 (0.32)	1	0.06 (0.23)
# Water birds	4	0.19 (0.45)	3	0.11 (0.31)	5	0.42 (1.16)	8	0.69 (2.36)	3	0.22 (0.76)	2	0.14 (0.68)

Table D-2 (cont'd).—Total, mean, and standard deviation by species guilds for the Burned Area from 2011 to 2016.

Los Lunas Burned area	2011 12 points		2012 12 points		2013 12 points		2014 12 points		2015 12 points		2016 12 points	
	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)	Total	Mean (SD)
# Species	30	5.44 (0.81)	30	5.72 (0.74)	23	5.61 (0.87)	28	5.50 (1.21)	32	4.97 (1.11)	38	5.14 (1.73)
# Birds	96	8.03 (2.08)	96	7.97 (1.73)	102	8.53 (2.08)	100	8.31 (2.27)	73	6.06 (1.96)	91	7.58 (1.13)
# Canopy spp.	2	0.11 (0.32)	3	0.17 (0.38)	3	0.19 (0.47)	5	0.25 (0.50)	4	0.25 (0.50)	4	0.22 (0.42)
# Canopy birds	2	0.11 (0.32)	2	0.17 (0.38)	3	0.25 (0.65)	5	0.28 (0.57)	4	0.28 (0.57)	4	0.31 (0.67)
# Cavity spp.	3	0.33 (0.53)	6	0.58 (0.60)	4	0.39 (0.55)	4	0.22 (0.48)	7	0.44 (0.69)	7	0.58 (0.60)
# Cavity birds	5	0.47 (0.91)	9	0.78 (0.90)	6	0.47 (0.70)	4	0.25 (0.55)	7	0.53 (0.88)	10	0.86 (1.25)
# Dense shrub spp.	2	0.19 (0.40)	2	0.14 (0.35)	0	0.00 (0.00)	2	0.22 (0.42)	1	0.31 (0.47)	2	0.17 (0.38)
# Dense shrub birds	3	0.22 (0.48)	2	0.19 (0.52)	0	0.00 (0.00)	3	0.25 (0.50)	4	0.31 (0.47)	2	0.17 (0.38)
# Edge spp.	3	0.92 (0.44)	4	0.92 (0.50)	2	0.94 (0.33)	1	0.75 (0.44)	4	0.89 (0.52)	4	0.47 (0.51)
# Edge birds	13	1.08 (0.60)	15	1.25 (1.00)	18	1.53 (0.84)	16	1.31 (1.01)	13	1.11 (0.78)	9	0.72 (0.94)
# Ground shrub spp.	4	0.83 (0.61)	4	1.00 (0.72)	4	1.00 (0.72)	5	1.06 (0.75)	5	0.72 (0.78)	7	0.94 (0.92)
# Ground shrub birds	17	1.39 (1.23)	14	1.19 (0.89)	16	1.33 (1.10)	18	1.47 (1.23)	10	0.86 (0.93)	16	1.28 (1.23)
# Mid-story spp.	12	2.83 (1.06)	10	2.78 (1.05)	9	2.94 (0.98)	9	2.89 (0.98)	10	2.33 (1.24)	11	2.56 (1.38)
# Mid-story birds	50	4.19 (1.83)	48	3.97 (1.76)	56	4.67 (2.01)	55	4.58 (1.79)	35	2.94 (2.11)	48	3.97 (2.57)
# Opening spp.	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)
# Opening birds	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)	0	0.00 (0.00)
# Water spp.	4	0.22 (0.48)	1	0.14 (0.35)	1	0.14 (0.35)	2	0.11 (0.40)	1	0.03 (0.17)	4	0.17 (0.45)
# Water birds	7	0.56 (1.52)	4	0.33 (0.93)	3	0.28 (0.74)	2	0.17 (0.61)	1	0.03 (0.17)	4	0.25 (0.69)

Appendix E

Southwestern Willow Flycatcher Survey Forms and Maps
and
Western Yellow-billed Cuckoo Survey Forms
2016

Willow Flycatcher (WIFL) Survey and Detection Form (revised April, 2010)

Site Name: BL-25 State: New Mexico County: Valencia
 USGS Quad Name: Toime, Los Lunas Elevation: 1,469 (meters)
 Creek, River, or Lake Name: Rio Grande

Is copy of USGS map marked with survey area and WIFL sightings attached (as required)? Yes X No
 Survey Coordinates: Start: E 341,191 N 3,848,584 UTM Datum: NAD83 (See instructions)
 Stop: E 340,201 N 3,845,501 UTM Zone: 13

If survey coordinates changed between visits, enter coordinates for each survey in comments section on back of this page.

****Fill in additional site information on back of this page****

Survey # Observer(s) (Full Name)	Date (m/d/y) Survey Time	Number of Adult WIFLs	Estimated Number of Pairs	Estimated Number of Territories	Nest(s) Found? Y or N If Yes, number of nests	Comments (e.g., bird behavior, evidence of pairs or breeding-potential threats [livestock, cowbirds, <i>Diochabes</i> spp.]). If <i>Diochabes</i> found, contact USFWS and State WIFL coordinator.	GPS Coordinates for WIFL Detections (this is an optional column for documenting individuals, pairs, or groups of birds found on each survey). Include additional sheets if necessary.			
Survey # 1 Observer(s): A. Cressotti	Date: 5/24/2016	0	0	0	N	Site dry with dense vegetation throughout. Cowbirds detected.	# Birds	Sex	UTME	UTMN
	Start: 5:45									
	Stop: 10:45									
	Total hrs: 5.0									
Survey # 2 Observer(s): J. Tinas	Date: 6/23/2016	0	0	0	N	Site dry with dense understorey and mature overstorey canopy.	# Birds	Sex	UTME	UTMN
	Start: 6:30									
	Stop: 11:00									
	Total hrs: 4.5									
Survey # 3 Observer(s): H. Dzwang	Date: 7/12/2016	0	0	0	N	Large, dense willow patch in the middle of the site along the river. Site dry. Cowbirds and livestock detected.	# Birds	Sex	UTME	UTMN
	Start: 6:00									
	Stop: 10:00									
	Total hrs: 4.0									
Survey # 4 Observer(s):	Date:						# Birds	Sex	UTME	UTMN
	Start:									
	Stop:									
	Total hrs:									
Survey # 5 Observer(s):	Date:						# Birds	Sex	UTME	UTMN
	Start:									
	Stop:									
	Total hrs:									
Overall Site Summary <small>Totals do not equal the sum of each column. Include only resident adults. Do not include migrants, nestlings, and fledglings. Be careful not to double count individuals.</small>		Total Adult Residents	Total Pairs	Total Territories	Total Nests	Were any WIFLs color-banded? Yes <u> </u> No <u>X</u> If yes, report color combination(s) in the comments section on back of form and report to USFWS.				
Total survey hrs: 13.5		0	0	0	0					

Reporting Individual: Darrell Ahlers Date Report Completed: 9/7/2016
 US Fish & Wildlife Service Permit #: TE819475-5 State Wildlife Agency Permit #: N/A

Submit form to USFWS and State Wildlife Agency by September 1st. Retain a copy for your records.

Fill in the following information completely. Submit form by September 1st. Retain a copy for your records.

Reporting Individual Darrell Ahlers Phone # (303) 445-2233
 Affiliation Bureau of Reclamation E-mail dahlers@usbr.gov
 Site Name BL-25 Date report Completed 9/7/2016
 Was this site surveyed in a previous year? Yes X No Unknown
 Did you verify that this site name is consistent with that used in previous yrs? Yes X No Not Applicable
 If name is different, what name(s) was used in the past? N/A
 If site was surveyed last year, did you survey the same general area this year? Yes X No If no, summarize below.
 Did you survey the same general area during each visit to this site this year? Yes X No If no, summarize below.
 Management Authority for Survey Area: Federal Municipal/County State Tribal Private X
 Name of Management Entity or Owner (e.g., Tonto National Forest) MRGCD
 Length of area surveyed: 3.3 (km)

Vegetation Characteristics: Check (only one) category that best describes the predominant tree/shrub foliar layer at this site:

 Native broadleaf plants (entirely or almost entirely, > 90% native)
X Mixed native and exotic plants (mostly native, 50 - 90% native)
 Mixed native and exotic plants (mostly exotic, 50 - 90% exotic)
 Exotic/introduced plants (entirely or almost entirely, > 90% exotic)

Identify the 2-3 predominant tree/shrub species in order of dominance. Use scientific name.

Salix exigua, Eleagnus angustifolia, Populus sp.

Average height of canopy (Do not include a range): 15 (meters)

Attach the following: 1) copy of USGS quad/topographical map (REQUIRED) of survey area, outlining survey site and location of WIFL detections;

2) sketch or aerial photo showing site location, patch shape, survey route, location of any detected WIFLs or their nests;

3) photos of the interior of the patch, exterior of the patch, and overall site. Describe any unique habitat features in Comments.


Comments (such as start and end coordinates of survey area if changed among surveys, supplemental visits to sites, unique habitat features).

Attach additional sheets if necessary.

Territory Summary Table. Provide the following information for each verified territory at your site.

Territory Number	All Dates Detected	UTM E	UTM N	Pair Confirmed? Y or N	Nest Found? Y or N	Description of How You Confirmed Territory and Breeding Status (e.g., vocalization type, pair interactions, nesting attempts, behavior)

Attach additional sheets if necessary



Yellow-billed Cuckoo Survey Form														
Site Name: BL-25		County: Socorro				State: New Mexico								
USGS Quad Name: Tome						Elevation: 1,469m								
Creek, River, Wetland, or Lake Name: Rio Grande														
Site Coordinates: Start: E 341,191 N 3,848,584		UTM Zone: 13												
Stop: E 340,201 N 3,845,501		Datum: NAD83												
Ownership: Private														
Was site surveyed in previous year?		<div style="display: flex; justify-content: space-between;"> Yes If yes, what site name was used? BL-25 </div>												
Survey # Observer(s) (Last Name, First Initial)	Date (m/d/y) Time, Total Hours	Total Number of YBCUs detected	Time Detected (AM)	Detect Type: I-Incidental P-Playback A-aerial V-visual B-both	Voc. Type: CN-Contact CO-coo AL-alarm OT-other (describe)	Playback #: Number of times Kowdy call played prior to response	Behavior code	Surveyor Detection Coordinates		Distance (m)	Bearing	Cuckoo #	Corrected Coordinates	
								UTM E	UTM N				UTM E	UTM N
Survey Period #1 Observer(s): Tinges	Date:													
	6/23/2016													
	Start:													
	8:30 AM													
	Stop:													
	11:00 AM													
	Total hrs:													
	4:50	0												
Survey Period #2 Observer(s): Dreong	Date:													
	7/12/2016													
	Start:													
	6:00 AM													
	Stop:													
	10:00 AM													
	Total hrs:													
	4:00	0												
Survey Period #3 Observer(s): Suche	Date:													
	7/25/2016													
	Start:													
	6:00 AM													
	Stop:													
	10:30 AM													
	Total hrs:													
	4:50	0												
Survey Period #4 Observer(s): Crescenti	Date:													
	8/8/2016													
	Start:													
	6:15 AM													
	Stop:													
	10:15 AM													
	Total hrs:													
	4:00	0												
Survey Period #5 Observer(s):	Date:													
	Start:													
	Stop:													
	Total hrs:													
Survey Summary:		# Det	# PO	# PR	# CO	# Nests found	Total Survey Hours:							
Total YBCUs*		0	0	0	0	0	17.00							
Notes (refer to Cuckoo # associated with individual detections)		Majority of habitat was intermediate to mature mixed canopy												
		Cottonwood overstory throughout site with sparse coyote willow and Russian olive understory												
		Willows near river and cottonwood overstory through most of the site, adequate habitat for cuckoo's												
		Cottonwood canopy runs the length of the site moderate habitat in areas												
*Include justification for these designations.														
Behavior Codes: AN = at nest, BI = brooding or incubating, CF = adult carrying food, CN = carrying nest material, COP = copulation, CP = catches prey, DD = distraction displays/defense of nesting area, EF = eats food, FL = recently fledged young of species incapable of flight, FLY = flying, FO = foraging, FS = adult carrying a fecal sac, FY = adult feeding nestlings, JUV = juvenile, NB = nest building, NE = active nest with unbroken eggs in it, NY = nest with young seen or heard in it, ON = occupied nest, PR = preening, SI = sitting, UIS = used, inactive nest with blue-green eggshells.														

Appendix F

Plant list
and
Total Percent Cover of Plants Detected in the Understory Layer
by Individual Species, Life-form, and Cover Type
2003 to 2016

Table F-1.—Plant list of species detected from 2003 to 2016.

	Code	Scientific name	Common name	Lifeform
Trees/shrubs	BASA	<i>Baccharis salicifolia</i>	Seep willow	NS
	ELAN	<i>Eleagnus angustifolia</i>	Russian olive	IT
	POAN	<i>Populus angustifolia</i>	Narrowleaf cottonwood	NT
	PODE	<i>Populus deltoides</i>	Rio Grande cottonwood	NT
	SAEX	<i>Salix exigua</i>	Coyote willow	NT/S
	SAGO	<i>Salix gooddingii</i>	Gooddings willow	NT
	TARA	<i>Tamarix ramosissima</i>	Saltcedar	IT/S
	ULPU	<i>Ulmus pumila</i>	Siberian elm	IT
Grasses/grass-like	AGGI	<i>Agrostis gigantea</i>	Redtop	IG
	BOBA	<i>Bothriochloa barbinodis</i>	Cane bluestem	NG
	BRIN	<i>Bromus inermis</i>	Smooth brome	IG
	BRJA	<i>Bromus japonicus</i>	Japanese brome	IG
	CASP	<i>Carex</i> sp.	Sedge	NG
	COSE	<i>Cortaderia selloana</i>	Pampas grass	IG
	CYOD	<i>Cyperus odoratus</i>	Fragrant flatsedge	NG
	DISP	<i>Distichlis spicata</i>	Saltgrass	NG
	ECCR	<i>Echinochloa crus-galli</i>	Barnyard grass	IG
	ELPA	<i>Eleocharis palustris</i>	Common spikerush	NG
	ELCA	<i>Elymus canadensis</i>	Canada wildrye	NG
	ELTR	<i>Elymus trachycaulus</i>	Slender wheatgrass	NG
	ERHY	<i>Eragrostis hypnoides</i>	Teal lovegrass	NG
	HOJU	<i>Hordeum jubatum</i>	Barley foxtail	NG
	JUBA	<i>Juncus balticus</i>	Baltic rush	NG
	JUEN	<i>Juncus ensifolius</i>	Sword-leaved rush	NG
	LEOR	<i>Leersia oryzoides</i>	Rice cutgrass	NG
	LEFU	<i>Leptochloa fusca</i>	Mexican sprangletop	NG
	MUAS	<i>Muhlenbergia asperifolia</i>	Scratchgrass	NG
	MURA	<i>Muhlenbergia racemosa</i>	Muhly	NG
	PACA	<i>Panicum capillare</i>	Witchgrass	NG
	PAOB	<i>Panicum obtusum</i>	Vine mesquite	NG
	PHAU	<i>Phragmites australis</i>	Common reed	NG
	POPA	<i>Poa pratensis</i>	Kentucky bluegrass	NG
	POMO	<i>Polypogon monspeliensis</i>	Rabbitfoot grass	IG
	SCPR	<i>Schedonorus pratensis</i>	Meadow fescue	IG
	SCAC	<i>Schoenoplectus acutis</i>	Hardstem bulrush	NG
	SCAM	<i>Schoenoplectus americanus</i>	American threesquare	NG
	SPAI	<i>Sporobolus airoides</i>	Alkali sacaton	NG
	SPCR	<i>Sporobolus cryptandrus</i>	Sand dropseed	NG
Forbs	AGPA	<i>Agastache pallidiflora</i> ssp <i>neomexicana</i>	New Mexico giant hyssop	NF
	AMBL	<i>Amaranthus blitoides</i>	Prostrate amaranth	IF
	AMPS	<i>Ambrosia psilostachya</i>	Western ragweed	NF
	APCA	<i>Apocynum cannabinum</i>	Clasping-leaf dogbane	NF
	ARAB	<i>Artemisia absinthium</i>	Wormwood	IF
	ARAN	<i>Argentina anserina</i>	Silverweed cinquefoil	NF
	ASSU	<i>Asclepias subverticillata</i>	Horsetail milkweed	NF
	ASSP	<i>Astragalus</i> sp.	Milkvetch	NF
	BIFR	<i>Bidens frondosa</i>	Beggarstick	NF
	CHAL	<i>Chenopodium album</i>	Lambsquarters	IF
	CHSE	<i>Chamaesyce serpyllifolia</i>	Thymeleaf spurge	NF
	CLLI	<i>Clematis ligusticifolia</i>	Virgin's bower	NF
	COAR	<i>Convolvulus arvensis</i>	Field bindweed	IF
	COCA	<i>Conyza canadensis</i>	Horseweed	NF
	CUSP	<i>Cuscuta</i> sp.	Dodder	NF
	DALE	<i>Dalea leporina</i>	Foxtail dalea	NF

	Code	Scientific name	Common name	Lifeform
	DEIL	<i>Desmanthus illinoensis</i>	Bundleflower	NF
	EQLA	<i>Equisetum laevigatum</i>	Smooth scouringrush	NF
	EUOC	<i>Euthamia occidentalis</i>	Western goldentop	NF
	GAPA	<i>Gaura parviflora</i>	Small-flowered gaura	NF
	GRSQ	<i>Grindelia squarrosa</i>	Curlycup gumweed	NF
	HEAN	<i>Helianthus annuus</i>	Common sunflower	NF
	KOSC	<i>Kochia scoparia</i>	Kochia	IF
	LASP	<i>Lactuca serriola</i>	Prickly lettuce	IF
	LELA	<i>Lepidium latifolium</i>	Perennial pepperweed	IF
	MEAL	<i>Melilotus albus</i>	White sweetclover	IF
	OEEL	<i>Oenothera elata</i>	Hooker's evening primrose	NF
	PESP	<i>Penstemon</i> sp.	Penstemon	NF
	PLLA	<i>Plantago lanceolata</i>	Narrowleaf plantain	IF
	PLMA	<i>Plantago major</i>	Common plantain	IF
	POLA	<i>Polygonum lapathifolium</i>	Pale smartweed	NF
	PSST	<i>Pseudognaphalium stramineum</i>	Cottonbatting cudweed	NF
	PYPa	<i>Pyrrhopappus pauciflorus</i>	Smallflower desert-chicory	NF
	RATA	<i>Ratibida tagetes</i>	Short-rayed coneflower	NF
	RUCR	<i>Rumex crispis</i>	Curly dock	IF
	SAIB	<i>Salsola iberica</i>	Russian thistle	IF
	SOAR	<i>Sonchus arvensis</i>	Field sowthistle	IF
	SOCA	<i>Solidago canadensis</i>	Golden rod	NF
	SYER	<i>Symphyotrichum ericoides</i>	White heath aster	NF
	TAOF	<i>Taraxacum officinale</i>	Dandelion	IF
	TRTE	<i>Tribulus terrestris</i>	Goats head	IF
	XAST	<i>Xanthium strumarium</i>	Common cocklebur	NF

*NT/S=Native tree/shrub; IT/S=Introduced tree/shrub; NG=Native grass; IG=Introduced grass; NF=Native forb; IF=Introduced forb

Table F-2.— Total percent cover of by individual species, life-form and cover type in the understory layer.

Understory layer	Total Percent Cover													
	2003	2004	2005	2006	2007*	2008	2009	2010	2011	2012	2013	2014	2015	2016
Coyote willow	0.6	1.0	1.9	4.7	0.5	1.0	1.3	1.1	2.2	0.8	3.4	1.9	0.9	1.1
Cottonwood	0.0	0.4	1.3	7.1	0.3	0.5	0.3	0.1	0.4	0.3	0.5	0.1	0.0	0.0
Gooddings willow	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0
Total native shrubs	0.6	1.4	3.2	11.9	0.9	1.5	1.7	1.2	2.7	1.1	3.9	2.0	0.9	1.1
Saltcedar	0.4	0.8	2.8	5.0	1.0	0.8	0.8	1.1	1.3	0.7	1.1	1.1	0.6	0.8
Russian olive	0.0	0.0	0.0	0.2	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
Siberian elm	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2
Total introduced shrubs	0.4	0.8	2.8	5.2	1.0	0.9	0.8	1.1	1.3	0.7	1.1	1.2	0.8	1.0
Fragrant flatsedge	1.7	3.5	8.4	0.5	2.1	4.4	1.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Baltic rush	1.3	0.0	0.0	0.0	0.0	0.0	1.1	0.7	0.3	0.2	0.0	0.1	0.3	0.1
Muhly	1.3	2.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Witchgrass	1.1	5.2	4.4	0.8	0.4	1.7	0.4	0.4	0.3	0.0	0.3	0.0	0.2	0.0
Vine mesquite	0.4	0.4	1.6	4.7	7.6	12.2	16.9	15.7	9.2	4.5	6.7	6.7	9.2	5.7
Common spikerush	0.0	0.2	0.0	0.0	0.2	0.4	0.5	0.4	0.0	0.0	0.0	0.0	0.0	0.0
Saltgrass	0.0	0.1	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	1.0	0.0	0.3	1.8
Kentucky bluegrass	0.0	0.2	0.6	0.3	0.1	0.0	0.0	0.4	0.0	0.1	0.0	0.4	0.4	0.0
Sedge	0.0	0.1	0.0	0.0	0.1	0.6	0.1	0.7	0.6	0.8	0.5	0.3	1.6	1.8
Mexican sprangletop	2.2	6.7	1.1	2.5	0.1	0.7	0.4	0.2	0.0	0.0	0.1	0.0	0.0	0.0
Teal lovegrass	0.0	0.0	2.6	0.0	0.3	0.2	0.2	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Barley foxtail	0.0	0.0	0.0	2.8	5.3	7.4	2.6	4.2	0.4	0.3	0.1	0.4	0.3	1.5
Common reed	0.0	0.0	0.0	0.0	0.8	0.4	0.6	0.7	0.7	1.0	0.5	0.3	0.4	0.3
Sword-leaved rush	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0
Rice cutgrass	0.0	0.0	0.0	0.0	0.0	0.2	0.5	0.2	0.0	0.0	0.0	0.0	0.0	0.0
Hardstem bulrush	0.0	0.0	0.0	0.0	0.0	0.4	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.3
American threesquare	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.2	0.1	0.0	0.0	0.0	0.0	0.0
Scratchgrass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.9	1.8	0.3	1.0	1.4
Sand dropseed	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.7	0.5	0.3	1.0
Slender wheatgrass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.2	0.0	0.0	0.0	0.3
Cane bluestem	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.1	0.3	0.0
Alkali sacaton	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.3	0.4	0.4
Canada wildrye	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
Total native grasses	8.0	19.1	18.7	11.6	17.0	28.8	25.4	24.7	12.4	8.1	12.4	9.4	14.7	14.7
Barnyard grass	1.3	4.3	6.0	2.8	1.0	1.1	0.4	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Rabbitfoot grass	1.6	4.5	2.8	0.1	2.0	3.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Smooth brome	0.0	0.0	0.0	0.0	0.0	1.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Meadow fescue	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.4	0.5	0.6	0.4	0.6	0.4	0.7
Japanese brome	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	0.3	0.2	0.0	1.2
Pampas grass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.2	0.2	0.0	0.3	0.1
Redtop	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4
Total introduced grasses	2.9	8.8	8.8	2.9	3.0	5.7	0.9	1.3	0.5	1.3	0.9	0.8	0.7	2.4
Horseweed	0.2	0.0	0.0	4.3	7.7	0.0	0.0	0.7	0.3	0.6	0.4	4.1	0.2	0
Common sunflower	7.9	13.9	0.3	3.9	1.1	1.9	0.0	1.0	0.0	0.8	0.2	0.3	0.0	0.0
Pale smartweed	0.8	1.2	0.2	5.9	1.0	0.2	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Common cocklebur	0.3	3.3	17.9	8.1	10.3	19.4	11.8	3.8	0.1	0.2	1.2	1.2	1.1	0.6
Beggarstick	0.0	0.9	3.4	0.5	0.1	0.2	0.2	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Western goldentop	0.0	0.3	0.8	1.7	2.9	11.9	9.2	7.3	3.4	2.8	2.3	2.6	3.9	3.9
Clasping-leaf dogbane	0.0	0.0	0.3	0.2	0.9	1.5	1.5	1.4	1.5	1.3	1.3	1.5	1.0	3.3
Milkvetch	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cottonbatting cudweed	0.0	0.0	0.0	1.2	0.6	0.0	0.2	0.1	0.3	0.0	0.0	0.0	0.0	0.0
Hooker's evening primrose	0.0	0.0	0.0	1.2	0.0	0.2	0.1	0.7	0.4	0.0	0.1	0.2	0.3	0.1
Dodder	0.0	0.0	0.0	0.1	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Bundleflower	0.0	0.0	0.0	0.0	0.5	0.2	0.7	1.3	0.2	0.0	0.0	0.0	0.0	0.0
Western ragweed	0.0	0.0	0.0	0.2	0.4	0.8	1.3	2.0	2.5	2.7	2.5	2.3	3.8	3.9
Silverweed cinquefoil	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Penstemon	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Smooth scouringrush	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.8	0.7	1.0	0.4	0.4	0.2	1.4
New Mexico giant hyssop	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.2	0.1	0.1	0.0	0.0	0.0	0.0
Curlycup gumweed	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Thymeleaf spurge	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.1	0.8	0.0	0.3	0.3
Small-flowered gaura	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.3	0.0	0.0
Foxtail dalea	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.1	0.3	0.0	0.0	0.0

Golden rod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.0	0.0
Short-rayed coneflower	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.1	0.3	0.0
Horsetail milkweed	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.3	0.9
Vigin's bower	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.1
White heath aster	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0
Smallflower desert-chicory	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3
Total native forbs	9.2	19.6	22.9	27.5	25.5	37.0	26.1	19.7	9.8	10.0	10.3	13.7	11.5	14.8
Lambsquarters	6.2	5.2	0.3	0.1	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Kochia	0.5	3.6	3.8	4.2	2.8	2.7	2.7	3.3	0.0	3.0	2.1	1.8	2.2	1.0
Prickly lettuce	0.1	0.8	0.0	6.0	2.3	0.9	0.0	0.2	0.1	0.6	0.1	0.2	0.0	0.1
White sweetclover	4.2	7.1	0.4	6.8	4.7	1.7	1.5	1.2	4.4	2.7	3.5	7.3	1.8	0.5
Russian thistle	0.0	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Perennial pepperweed	0.0	0.2	0.0	0.0	0.0	0.0	0.1	2.3	0.3	1.0	0.3	0.1	0.0	0.1
Wormwood	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Curly dock	0.0	0.0	0.1	0.5	1.6	0.1	0.0	0.1	0.3	0.0	0.1	0.0	0.0	0.0
Prostrate amaranth	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Goats head	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Field bindweed	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.5	0.3	0.2	0.1
Narrowleaf plantain	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0
Dandelion	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Common plantain	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0
Field sowthistle	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.1	0.1
Total Introduced forbs	11.0	17.8	4.8	17.8	11.4	5.7	4.4	7.0	5.3	7.3	6.7	10.0	4.3	1.9
Total understory vegetation	32.1	67.5	61.2	76.9	58.8	79.6	59.3	55.0	32.0	28.5	35.3	37.1	32.9	35.9
Litter	4.4	5.2	7.3	5.5	23.4	12.7	30.5	42.6	60.1	67.8	55.3	59.3	65.7	57.2
Bare soil	63.5	27.3	31.5	17.6	17.8	7.7	10.2	2.4	7.9	3.7	9.4	3.7	1.4	7.0
Total cover	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.1	100.0	100.1

Appendix G

Groundwater Monitoring Wells
Monthly Data
June 2003 – October 2010

Table G-1.—Depth (in inches) below the ground surface to water at each well for each monthly reading from June 2004 to October 2010.

Date	Well number (depth of well)										
	N1 (62)	N2 (62)	N3 (60.5)	N4 (64)	M1 (59)	M2 (61)	M3 (59)	M4 (61)	S1 (56)	S2 (61.5)	S3 (69)
06/04/03	44.0	41.0	29.0	No well	30.0	29.0	28.0	No well	34.0	49.0	No well
09/04/03	dry	dry	dry	No well	dry	dry	dry	No well	dry	dry	No well
10/30/03	45.0	41.0	31.0	No well	32.0	32.5	36.5	No well	40.0	dry	No well
11/27/03	36.0	41.0	37.0	No well	20.0	19.0	22.5	No well	28.5	51.0	No well
12/21/03	37.0	33.0	25.0	No well	20.0	20.0	21.5	No well	30.5	53.0	No well
01/24/04	38.0	33.0	23.0	No well	20.5	19.5	20.5	No well	31.0	53.0	No well
03/11/04	38.5	33.5	23.5	No well	21.5	20.5	20.5	No well	32.0	54.0	No well
04/01/04	32.0	27.5	18.5	No well	15.5	15.5	18.0	No well	27.5	50.5	No well
04/30/04	42.0	37.0	26.0	No well	26.5	25.5	25.5	No well	37.5	60.0	No well
05/30/04	35.5	33.0	24.0	No well	19.5	20.5	21.5	No well	31.5	55.5	No well
06/29/04	53.5	47.5	35.0	No well	39.5	37.0	36.5	No well	48.5	dry	No well
08/05/04	57.0	53.0	46.0	42.0	31.0	41.0	41.5	dry	39.5	dry	65.0
09/02/04	dry	dry	dry	58.0	dry	dry	dry	dry	56.0	dry	66.0
10/05/04	54.0	49.0	37.0	39.5	41.5	42.0	46.5	dry	50.5	dry	64.0
11/05/04	42.0	37.0	26.0	31.0	28.0	No well	29.5	41.0	35.5	58.0	49.0
12/04/04	36.5	30.0	19.0	23.5	20.0	No well	17.5	28.0	27.5	48.5	41.0
01/07/05	36.5	32.0	23.5	30.0	19.0	20.0	21.0	36.5	29.5	51.0	45.0
02/04/05	36.5	32.0	23.0	29.5	19.0	16.0	20.0	34.5	29.5	51.0	44.0
03/03/05	30.0	27.0	19.0	27.5	13.0	11.0	16.0	33.0	23.0	45.5	39.5
04/02/05	26.5	24.0	16.0	26.0	10.0	8.5	13.0	32.0	19.0	42.0	37.0
05/06/05	0.0	14.5	8.5	19.0	0.0	0.0	5.5	25.5	11.0	36.0	32.5
06/06/05	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
07/31/05	dry	57.5	43.0	40.5	47.0	39.5	42.0	49.5	52.0	dry	61.5
08/30/05	dry	59.0	40.0	34.0	48.0	40.0	37.5	52.0	52.5	dry	63.0
09/30/05	56.0	47.0	34.0	35.5	26.0	26.0	34.5	47.0	39.5	dry	56.0
10/31/05	52.0	43.5	31.0	34.0	28.0	24.5	29.0	43.5	34.5	56.5	48.5
11/29/05	45.5	38.0	27.0	32.0	22.5	20.0	25.0	40.0	30.0	52.0	45.5
12/30/05	42.5	35.0	23.5	28.0	21.0	17.0	21.5	33.0	29.0	50.0	43.5
01/31/06	46.5	39.0	27.5	32.5	24.0	21.0	25.0	38.0	34.0	54.5	46.5
02/28/06	48.0	40.0	28.5	32.5	26.5	22.5	25.0	38.5	36.5	56.5	49.0
03/31/06	59.5	49.5	35.0	36.0	39.5	32.5	34.5	44.5	46.0	dry	55.5
04/28/06	57.5	48.5	36.0	37.0	38.0	32.0	35.5	47.0	43.0	dry	54.5
05/29/06	53.5	46.5	36.0	38.0	32.0	29.0	34.5	47.5	39.0	dry	53.0
06/30/06	54.0	45.0	32.0	33.5	37.0	31.0	33.0	42.5	40.5	60.0	50.0
07/26/06	dry	55.0	39.5	36.0	52.0	43.5	43.5	49.0	55.5	dry	60.5
08/28/06	55.5	46.5	33.0	33.5	39.0	32.5	33.5	43.0	42.0	dry	52.5
09/21/06	dry	53.5	38.5	38.0	48.0	40.0	41.5	50.0	52.0	dry	60.5
10/31/06	42.0	35.0	36.0	29.5	19.0	17.0	22.5	36.5	26.5	49.5	43.0
11/30/06	41.5	36.0	29.5	24.5	15.0	13.0	17.5	33.0	23.5	46.5	40.5
01/27/07	43.5	36.5	26.0	31.5	21.5	18.5	22.0	36.5	31.5	53.0	45.5
02/26/07	43.0	36.0	25.5	31.0	21.0	18.0	21.5	36.0	31.0	52.5	45.0
03/28/07	29.0	24.0	15.0	22.5	9.5	7.5	12.0	28.0	20.0	42.0	36.0
04/29/07	46.5	37.5	25.5	28.5	29.5	24.0	26.0	37.5	36.0	56.5	47.0
05/31/07	27.5	21.5	17.5	25.0	10.5	9.5	14.5	32.5	20.0	56.5	38.0

Date	Well number (depth of well)										
	N1 (62)	N2 (62)	N3 (60.5)	N4 (64)	M1 (59)	M2 (61)	M3 (59)	M4 (61)	S1 (56)	S2 (61.5)	S3 (69)
06/29/07	50.0	41.5	28.0	29.0	37.5	32.5	34.5	43.0	42.5	dry	51.5
07/31/07	51.5	44.0	31.5	33.0	36.5	32.0	35.5	46.0	41.5	dry	53.5
08/31/07	56.0	47.0	33.0	31.0	42.0	36.0	38.5	45.5	47.0	dry	54.0
09/28/07	57.5	47.0	34.5	35.0	42.5	36.5	38.5	47.5	47.5	dry	56.5
10/30/07	51.0	44.0	31.0	34.5	34.0	33.0	39.5	50.0	43.0	dry	54.5
11/30/07	46.5	40.5	29.0	33.5	30.5	30.5	33.5	46.5	38.5	58.0	51.5
12/28/07	40.0	34.0	25.0	30.5	22.5	19.0	22.5	37.5	31.5	53.0	46.0
01/29/08	37.5	32.5	23.0	29.5	19.5	17.5	22.0	37.5	29.5	51.5	44.5
02/29/08	29.0	26.0	18.0	26.0	11.0	10.0	16.0	33.0	20.5	43.0	38.0
03/31/08	17.0	14.0	6.0	15.0	1.0	0.0	6.5	22.0	9.5	33.0	28.0
04/28/08	14.0	10.5	3.5	14.0	-4.0	-2.5	5.0	21.5	6.5	30.5	26.0
05/28/08	12.0	12.0	2.0	13.5	-5.0	-3.5	4.5	21.5	5.5	32.0	26.5
06/30/08	35.0	30.0	19.0	22.0	24.0	18.5	10.0	31.5	28.5	50.5	40.5
07/28/08	49.0	41.5	28.0	28.5	36.0	29.5	32.0	38.5	40.0	dry	51.5
08/27/08	59.0	49.0	34.0	35.0	42.0	36.0	37.5	46.0	45.5	dry	55.0
09/27/08	58.0	48.0	32.5	32.0	41.0	34.5	36.5	44.0	45.5	dry	56.0
10/31/09	52.5	44.0	30.0	32.5	33.5	28.5	32.0	42.5	39.5	dry	51.5
11/29/08	43.0	36.5	25.5	30.0	28.0	23.5	26.5	39.0	34.5	56.5	48.0
12/30/08	43.0	36.0	25.0	29.5	25.5	22.0	25.5	38.0	33.5	55.5	47.5
01/31/09	43.5	36.0	25.0	29.5	26.0	22.0	25.0	38.0	33.5	55.0	47.0
02/28/09	38.0	31.0	19.0	22.5	23.0	18.5	22.5	34.0	31.0	52.0	44.5
03/30/09	35.0	28.5	17.0	21.0	19.5	16.0	21.0	33.0	28.0	50.0	42.0
04/27/09	19.0	17.5	10.0	17.5	1.5	2.0	10.5	25.5	9.5	35.5	29.5
05/25/09	6.5	17.0	8.0	17.0	-0.5	0.5	6.5	23.5	9.0	34.5	30.0
07/02/09	36.0	32.0	19.5	24.5	24.0	20.5	25.0	37.0	35.1	50.5	42.0
09/07/09	dry	dry	36.0	34.5	45.5	38.0	39.5	47.5	44.5	dry	52.5
10/09/09	dry	dry	37.0	36.0	46.5	38.5	40.0	47.5	45.5	dry	54.0
11/02/09	55.5	45.0	31.5	32.5	35.0	29.0	32.0	41.5	37.5	58.5	49.0
12/02/09	50.5	42.0	30.0	33.5	27.5	23.0	26.5	39.5	31.5	53.5	44.5
01/04/10	48.5	40.5	29.5	33.5	26.5	22.5	26.0	40.0	32.0	53.0	44.0
02/08/10	45.0	38.0	27.0	31.5	25.0	21.5	25.0	39.0	32.0	52.5	44.0
03/05/10	46.5	38.0	27.0	30.5	26.0	22.0	24.5	38.0	32.0	52.0	43.0
04/05/10	38.5	31.0	20.5	24.5	22.5	18.5	22.0	33.0	30.0	50.0	41.5
05/03/10	27.0	22.5	17.5	22.5	10.0	10.5	13.5	29.5	20.5	42.0	36.0
05/30/10	24.5	19.0	13.5	18.5	10.0	9.0	13.5	32.0	17.5	42.0	35.5
06/30/10	56.0	46.0	32.5	32.0	41.5	36.0	38.5	46.5	41.0	dry	51.0
07/31/10	49.0	41.5	30.0	31.0	33.0	29.0	33.5	44.0	35.0	58.0	47.5
08/30/10	dry	dry	41.0	dry	54.5	45.0	45.5	48.0	dry	dry	62.0
9/22/2010	dry	dry	50.0	43.0	dry	60.0	57.5	58.0	dry	dry	dry








Figure G-1.—Discharge (cfs) of the Rio Grande at San Acacia, New Mexico, and average ground water levels (inches from the surface) in wells along the South, Middle, and North transects at the LLRS, June 2003 to Oct. 2010.

Appendix H






Groundwater Monitoring Wells
HOBO Water Level Logger Data
June 2012 – September 2016

South Transect






Well S1 (Missing data from 9/15 to 9/16)



Well S2







Well S3








Middle Transect






Well M1 (Missing data from 9/15 to 9/16)



Well M2






Well M3 (Missing data 12/15 to 9/16)







North Transect





Well N1 (Missing data from 9/12 to 9/14)



Well N2 (Missing data from 9/15 to 9/16)



Well N3 (Missing data from 9/15 to 9/16)



Appendix I

Photo Stations
2003 - 2016

Photo Station 1 - Facing North



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 1 – Facing River



2003

No photo



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 1 – Facing South



2003



2004



2005



2006



2007



2008



2009




2010



2011



2012




2013



2014



2015



2016

Photo Station 2 – Facing North



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 2 – Facing River



2003

No photo



2005



2006



2007



2008



2009



2010



2011



2012




2013



2014



2015



2016

Photo Station 2 – Facing South



2003




2004



2005



2006



2007



2008



2009




2010



2011



2012




2013



2014



2015



2016

Photo Station 3 – Facing North



2003



2004



2005



2006



2007



2008



2009




2010



2011



2012



2013



2014



2015



2016

Photo Station 3 - Facing South



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 4 – Facing North



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 4 – Facing South



2003



2004



2005



2006



2007




2008



2009




2010



2011



2012



2013



2014



2015



2016

Photo Station 5 – Facing North



2003



2004



2005



2006



2007



2008




2009




2010



2011



2012



2013



2014



2015



2016

Photo Station 5 – Facing South



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 6 – Facing North

2003 No photo



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 6 – Facing South



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 7 – Facing North



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 8 – Pond



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 9 – Facing South



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

Photo Station 10 – Facing North



2003



2004



2005



2006



2007



2008



2009



2010



2011



2012



2013



2014



2015



2016

PEER REVIEW DOCUMENTATION

PROJECT AND DOCUMENT INFORMATION

Project Name Los Lunas Habitat Restoration Monitoring WOJD QA635

Document 2016 Monitoring Report for the Los Lunas Habitat Restoration Project

Document Date March 2017

Team Leader Darrell Ahlers

Document Author(s)/Preparer(s) Rebecca Siegle, Darrell Ahlers

Peer Reviewer Meghan White

Peer Reviewer _____

REVIEW REQUIREMENT

Part A: Document Does Not Require Peer Review

Explain _____

Part B: Document Requires Peer Review: SCOPE OF PEER REVIEW

Peer Review restricted to the following Items/Section(s): _____ Reviewer: _____

REVIEW CERTIFICATION

Peer Reviewer - I have reviewed the assigned Items/Section(s) noted for the above document and believe them to be in accordance with the project requirements, standards of the profession, and Reclamation policy.

Reviewer: Meghan White Review Date: March 2016 Signature: 

Reviewer: _____ Review Date: _____ Signature: _____

I have discussed the above document and review requirements with the Peer Reviewer and believe that this review is completed, and that the document will meet the requirements of the project.

Team Leader: Darrell Ahlers Date: 3/28/2017 Signature: 